CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 103, 2010

A

Abdel-Baky, N. F., 1909
Abdelghany, A. Y., 1909
Abel, C. A., 861
Abellán, J., 2164
Abernathy, R., 1886
Abney, M. R., 1268, 2087
Abolmaaty, A., 1547
Adamczyk, J., 1797
Addison, S. J., 1206
Afik, O., 228
Aguilar-Meza, O., 570
Ahmed, M. Z., 1848
Ahn, Y.-J., 1293
Akbar, W., 1431
Akotsen-Mensah, C., 744
Akratanakul, P., 1012
Alexander, C. E., 1896
Ali, S. E., 1621
Aliakbarpour, H., 631
Allemand, R., 308
Alm, S. R., 1574, 1810
Altmann, J., 2061
Aluja, M., 2000
Al-Zubaidy, M., 2042
An, J., 2169
Andow, D. A., 1394, 1803
Andrade, C.F.S., 166
Appel, A. G., 448, 762, 781, 2118
Arbogast, R. T., 991, 1002, 1503
Armstrong, J. S., 302, 356
Arredondo, J., 2000
Arthur, F. H., 197, 991, 1002
Aslam, M., 823
Athanassiou, C. G., 197, 977, 985
Atwa, A., 1621
Avilla, J., 482
Awadalla, S. S., 1909

B

Ba, N. M., 203
Babcock, J. M., 1031
Bacheler, J. S., 2236
Baker, G., 1347
Bambara, S. B., 1790
Bang, D. V., 127
Barak, A. V., 277, 603
Barbouletos, T., 242
Barbour, J. D., 641
Barrett, B. A., 1704
Barry, J. D., 284
Bartholomaeus, F., 1286
Bauer, L. S., 1080
Baur, M. E., 869
Beer, M.S.D., 1112
Beerli, P., 2214
Bello, G.M.D., 1094
Bello, N. M., 1242
Bennett, G. W., 172
Bernal, J., 1964
Bernal, J. L., 1964
Bernal, J. S., 570
Bernard, M. B., 2061
Bernklau, E. J., 2052
Bettiga, L. J., 272
Bi, J. L., 1172
Bielza, P., 2164
Bing, J. W., 1031
Binso-Dabire, C. L., 203
Bixby-Brosi, A. J., 1555
Bjostad, L. B., 2052
Blackburn, L. M., 1569
Blanco, C. A., 861
Blinka, E. L., 2236
Bo, L., 1988
Boase, C. J., 136
Boethel, D. J., 869
Boiteau, G., 63, 1613
Boll, R., 2242
Boozer, R., 744
Borgemeister, C., 1159
Bosch, D., 482
Bosque-Pérez, N. A., 691
Botha, A.-M., 887
Bouhssini, M. E., 958
Bourgeois, L., 917
Bourguet, D., 1803
Bout, A., 2242
Boykin, L. M., 2214
Bradley, J. R., 2236
Brazier, C., 1803
Brewer, M. J., 583
Brightwell, R. J., 1790
Brlansky, R. H., 1531
Broderick, N. A., 718
Broughton, S., 1744
Brown, B. D., 1569
Brown, J. K., 2174
Brown, M. W., 1657
Bruck, D. J., 416
Brunner, J. F., 140, 340, 1242, 1277, 1378
Bu, C. Y., 1372
Buchelos, C. T., 190
Buchholz, S., 1
Bueno, F. C., 676
Bueno, O. C., 676
Buitenhuis, R., 563
Bujang, N. S., 443
Bumgardner, M. S., 597
Buntin, G. D., 2072, 2229
Burd, J. D., 1454, 1875
Burgess, E.P.J., 497
Burkness, E. C., 1039
Buscarini, T., 708
Buschman, L. L., 348
Buss, E. A., 726
Butler, S. M., 1929
Byrne, F. J., 2174

C

Cañas, L. A., 949
Cabaleiro, C., 619
Caballero, P., 17, 577
Cagáň, L., 1803
Calvin, D. D., 249
Cambron, S. E., 2229
Campbell, J. F., 991, 1002
Cardona, C., 1860
Cardona, P. S., 938
Carinci, R., 973
Carriére, Y., 1821
Cartea, M. E., 1866
Cassanelli, S., 816
Castro, B. A., 1180
Chabi-Olaye, A., 1159
Chai, R.-Y., 460
Chambers, U., 416
Chareonviriyaphap, T., 1012
Chen, C., 2009
Chen, H., 1100
Chen, M., 2107
Chen, M.-S., 178
Chen, Yang, 1444
Chen, Yanping, 1517
Chen, Zhilin, 277
Chen, Zhinan, 277
Cheng, J., 1372
Chi, H., 1628
Chicca, M., 816
Chin, E., 172
Choe, D.-H., 1761
Choi, B.-R., 1293
Chong, K.-F., 1775
Chong, Y. V., 1994
Chouvenc, T., 1327
Christeller, J. T., 497
Chua, T. H., 1994
Chueca, P., 386, 1400
Cid, M., 619
Civolani, S., 816
Clark, T. L., 2187
Clarke, S. R., 95
Clough, G. H., 1306
Cognato, A. I., 1187, 1312
Cole, P., 2061
Coll, M., 1086

Colunga-Garcia, M., 108
Contreras, J., 2164
Cooper, W. R., 1875
Cornelius, M. L., 799
Corso, I. C., 869
Costamagna, A. C., 1483
Cottrell, T. E., 2072
Cowles, R. S., 1735, 1810
Cox, D. L., 770
Cox, R., 1797
Cox-Foster, D., 1517
Crespo, A.L.B., 1386
Crompton, D. S., 648
Crook, D. J., 1235
Crowe, C. M., 1216
Crull, L., 509
Currie, R. S., 348

D
Díaz-Fleischer, F., 2000
Da, K., 2072
Daane, K. M., 272, 1051, 1943
Dag, A., 228
Dammalage, T., 85
Davey, R. B., 211, 1360
David, N., 1306
Davidson, D., 219
Davila-Olivas, N. H., 949
Davis, H. N., 348
Davis, T. W., 119
Dean, D., 2214
DeBano, S. J., 1306
Decio, P., 676
Degain, B. A., 158, 2174
Delaney, K. J., 516
Delaplane, K. S., 1
Delay, C. E., 166
Delpuech, J.-M., 308
Dembilio, O., 402
Demuner, A. J., 1438
Dennehy, T. J., 158, 2174
Diaz, A., 2042
Diaz-Montano, J., 925
Diego, J. C., 1964
Dien, L. Q., 127
Dilling, C., 368
Diniz, A. N., 550
Diss-Torrance, A., 1569
Dodds, K. J., 698, 2094
Doerr, M. D., 140
Dogan, H., 1508
Dolatti, L., 958
Dorhout, D. L., 54
Downes, S., 472, 2147
Duan, J. J., 1080
Dubois, G. D., 698
Dupont, C., 308
Duque, J.E.L., 166
Dusfour, I., 973
Duvallet, G., 1012
Duyn, J. W. V., 2236

E
Ebert, T. A., 1531
Edwards, O., 958
Eigenbrode, S. D., 691
Eisenback, B. M., 1222
Ekhtelat, M., 2112
Ekobu, M., 1493
Elfékih, S., 1842
Ellersieck, M. R., 77, 2187
Ellis, J. D., 1
El-Rahman, S. A., 1621
El-Syrafi, H. A., 1909
Elvira, S., 17, 577
El-Zemaity, M. E.-S., 1547
Engels, H., 1803
Enríquez, C.-L. R., 662
Epsky, N. D., 533, 1886
Epstein, D., 1299
Esker, P. D., 1670
Espinosa, P. J., 2164
Espinoza, H. R., 1886
Eva, M. J., 2118
Evans, J. D., 1517
Evans, T. A., 423
Evenden, M. L., 654

F
Fabrick, J. A., 2174
Fadamiro, H. Y., 744
Fanhua, Z., 1988
Farrell, S. L., 1394
Fathipour, Y., 292, 1420, 2019
Ferguson, S., 2197
Fernández, E., 2164
Fernandes, F. L., 1438
Fields, P. G., 1909
Figueroa, J. I., 662
Fisher, T. W., 1060
Flanders, K. L., 2229
Flinn, P. W., 1508
Follett, P. A., 908
Fontana, J. D., 166
Foottit, R. G., 374
Fraguas, G. M., 166
Francese, J. A., 1235
Francisco, M., 1866
Fraser, I., 603, 1080, 1235
Frazier, J., 1517
Frazier, M., 1517
Frommer, M., 1071
Frost, K. E., 1670
Fuchs, M., 925
Fukumoto, T., 1784
Furlong, M., 2155
Fuse, C. B., 1094

G
Gaborit, P., 973
Gandhi, K.J.K., 1187
Gao, J., 2169
Gao, Y., 2169
Gao, Y.-Z., 2204
Garcerí, C., 386
Garcia, C. M., 101
Garczynski, S. F., 1060
Gardner, W. A., 754
Garrido-Bailón, E., 1964
Gassmann, A. J., 1821
Gautam, S. G., 1920
Gaylord, M. L., 1693
Gebre-Amlak, A., 302
Geden, C. J., 1937
George, J. E., 211
Gerry, A. C., 550
Gharalari, A. H., 219
Ghislain, M., 1493
Gibb, T., 172
Gilchrist, A. S., 1071
Giles, K. L., 1920
Gillani, W. A., 823
Ginzel, M. D., 147
Girod, R., 973
Glenn, D. M., 394
Goebel, P. C., 597
Goldfarb, H., 1299
Goldstein, J. A., 331
Golizadeh, A., 966
Gomez, L., 1594
González-Porto, A. V., 1964
Gorría, N., 17
Goto, C., 257
Gould, F., 861, 2169
Grávalos, C., 2164
Graaf, J. d., 1577
Grafton-Cardwell, E. E., 1400
Graham, E. E., 641
Grant, G. G., 1216
Grant, J., 368
Gratton, C., 1670
Greeff, J. M., 1848
Greene, J. K., 525
Gries, R., 654
Grieshop, M. J., 1242
Groot, A. T., 861
Grové, T., 1112
Guedes, R.N.C., 1438
Guo, S., 2028
Guo, W.-C., 1365
Guo, Y.-Y., 1444
Gut, L. J., 1299, 2100
Gwirtz, J. A., 1508

H
Haack, R. A., 108, 1682
Habibpour, B., 2112
Hahn, K.-W., 25
Hall, D. G., 541, 2214
Hallman, G. J., 34, 1129, 1950
Hamm, P. B., 1306
Hammon, R. W., 1875
Han, J., 1293
Han, Y., 1412
Hanavan, R. P., 691
Handelsman, J., 718

Hanks, L. M., 641
Hannon, E. R., 1821
Haraguchi, D., 1144, 2034
Harpold, V. S., 2174
Harris, M. K., 1312
Hartman, G. L., 509
Hashim-Buckey, J. M., 1051
Haubruge, E., 1517
Haverty, M. I., 770
Haymer, D. S., 1842
Heath, R. R., 533, 1886
Hegazi E. M., 1621
Helson, B., 708
Henderson, G., 1647, 2132
Heng-Moss, T. M., 1405
Henne, D. C., 1524
Hentz, M. G., 541
Herbert, A., 2236
Herman, T. K., 509
Herms, D. A., 597, 1187
Hernández, G., 861
Hibbard, B. E., 77, 2052, 2187
Higes, M., 1964
Hill, C. B., 509
Hinson, B., 219
Hirsch, J., 898
Hochmair, H. H., 1338
Hoddle, M. S., 877
Hodges, R. D., 1636
Hoebeke, E. R., 698
Hoffmann, E. J., 1196
Hofstetter, R. W., 1693
Hogendoorn, K., 1286
Holland, J. D., 1464, 2229
Holler, T., 1586
Hollingsworth, R. G., 908
Holmes, G. J., 2087
Hong, K.-J., 25
Hope, M., 1355
Horne, P. A., 2061
Horwood, M. A., 2140
Hou, M., 1412
Hoy, M. A., 726
Hu, C., 1444
Hu, L.-X., 1628
Hu, X., 1100
Hu, X. P., 762
Hu, Y., 1394
Hua, L., 1729
Huang, J., 2100
Huang, Y., 1832
Huckaba, R. M., 1031
Hunt, K. L., 1312
Hunt, T. E., 1405
Hussien, M. I., 1547
Hutchison, W. D., 1039

I

Iatrou, S. A., 190
Ibrahim, S. A., 2132
Ichinose, K., 127
Isaacs, R., 557, 668, 1720
Islam, A., 85
Issaly, J., 973

J

Jacas, J. A., 386, 402
Jager, A., 1508
James, W., 472
Jang, E. B., 1586, 1594
Jaramillo, J., 1159
Jenkins, D. A., 360
Jenkins, P. E., 1720
Jessup, A., 85
Jiang, W.-H., 1365
Jiao, Y., 277
Jiménez, J. J., 1964
Jin, Y. S., 1372
Johnson, J. A., 612
Johnson, S. A., 265
Jones, M. M., 1815
Jones, V. P., 1242
Jordan, T. A., 2080
Joseph, D., 2042
Joubert, P. H., 1112
Juárez, M. P., 1094
Judd, T. M., 2126
Jun, L., 95

K

Kalischuk-Tymensen, L. D., 1019
Kammlah, D. M., 211
Kang, L., 277
Kanga, L.H.B., 1797
Karanastasi, E., 977
Karimzadeh, J., 2019
Kaster, V., 2187
Kathuria, A., 2140
Kaufman, P. E., 1937
Kavallieratos, N. G., 190, 977
Keller, M. A., 1286
Kemp, D., 1355
Kemp, W. P., 234
Kendra, P. E., 533, 1886
Kennedy, G. G., 1268, 2087
Khafagi, W. E., 1621
Khamis, M., 1508
Khan, S. A., 938
Khanjani, M., 292
Khishen, A.A.E., 2187
Khocheili, F., 2112
Khrimian, A., 1586
Kim, S. I., 1293
Klocke, N. L., 348
Knutson, A. E., 735
Kobelt, A., 2061
Koehler, P. G., 1636
Kohama, T., 2034
Kok, L. T., 1222
Konstantopoulou, M. A., 1621
Krawczyk, G., 1657
Krupke, C. H., 147
Krupke, Christian H., 1464
Kuenen, L.P.S., 314
Kuenen, L.P.S. B., 1250
Kumano, N., 1144, 2034
Kuriwada, T., 1144, 2034
Kurtz, R., 2187
Kyamanywa, S., 1493

L

Lagalante, A. F., 1222
Lage, J., 958
Laine, R. A., 1647, 2132
Lambdin, P., 368
Lambkin, T. A., 2155
Lame, F.M.d., 1299
Lance, D. R., 1235
Landis, D. A., 1483
Laub, C. A., 2080
Lawrence, J., 1896
León, J.H.d., 908
Leathers, J., 2042
Leblanc, L., 409
Ledbetter, C., 1250
Lee, C.-Y., 437, 443, 460, 1754, 1770, 1775
Lee, K. V., 1039
Lee, R. D., 2072
Lee, S.-G., 1293
Leem, Y. J., 25
Lei, C., 158
Lei, Chaoliang, 2169
Leis, M., 816
Lema, M., 1866
Lemmetty, A., 1060
Leonard, B. R., 869
Leskey, T. C., 63
Leskey, Tracy C., 394
Levang-Brilz, N. M., 1950
Lewis, S. M., 1915
Lewis, V. R., 770
Lewis-Rosenblum, H., 1972
Li, A. Y., 1360
Li, B., 1100
Li, G., 1832
Li, G.-Q., 1365
Li, X., 158
Li, Xianchun, 2174
Li, Xiangyong, 843
Liao, W., 492
Liburd, O. E., 1258
Liesch, P. J., 1979
Lightle, D. M., 1187
Liu, Haijun, 1100
Liu, Hongbo, 492
Liu, Hui, 95
Liu, J., 1665
Liu, P., 1365
Liu, Tong-Xian, 1729
Liu, T.-X., 823
Liu, Xia, 1832
Liu, Xiang, 178, 958
Liu, Y.-B., 272
Liu, Y.-H., 2204
Llácer, E., 402

Lohmeyer, K. H., 211
Lord, J. C., 1542
Lowery, D. T., 374
Lu, W.-P., 1365
Lysyk, T. J., 1019

M

Ma, Jianyi, 492
Ma, Jun, 1100
Ma, W., 158
MacKinley, P. D., 1613
Maerere, A., 70
Mafra-Neto, A., 1594
Magalhães, S.T.V., 1438
Magarey, R. A., 108
Magsig-Castillo, J., 1172
Mahon, R. J., 472, 835, 2147
Maier, D. E., 1896
Mailleret, L., 2242
Makundi, R. H., 70
Malaspina, O., 676
Malinga, J., 958
Malone, L. A., 497
Malone, S., 2236
Manachini, B., 1803
Mandour, N. S., 1848
Mangini, A., 1216
Manicardi, G. C., 816
Mankin, R. W., 1135, 1636
Margosian, M. L., 108
Markin, G., 242
Markwick, N. P., 497
Marques, F. A., 166
Marques-da-Silva, E., 166
Marshall, J. L., 958
Marshall, K., 1797
Martínez, A.-M., 662
Martín-Hernández, R., 1964
Mashtoly, T. A., 1547
Mason, C. E., 331
Mason, L. J., 1896
Mastrangelo, T., 85
Mastro, V. C., 1235
Mathews, C. R., 1657
Mau, R.F.L., 409, 1594
McBrien, H. M., 1603
McCarthy, P. C., 1613
McCornack, B. P., 1483
McElfresh, J. S., 314, 1603
McFarlane, J., 708
McKenzie, N., 708
Meade, T., 1031
Meagher, R. L., 360
Meating, J., 708
Meihls, L. N., 77, 2187
Meinke, L. J., 849
Meloche, F., 63
Menti, H., 977
Menzies, M., 1355
Merrill, S. C., 302
Messenger, M., 603
Meyer, M. D., 70
Mian, M.A.R., 949
Michel, A. P., 949
Micoud, A., 1803
Midgarden, D., 1886
Milan, J., 242
Miles, J. W., 1860
Millar, J. G., 314, 641, 1216, 1603, 1943
Miller, D. R., 1216, 2094
Miller, J. R., 1299
Miller, R. J., 1360
Ming, Q.-L., 1915
Mitchell, R. F., 641
Miura, K., 1564
Miyatake, T., 1564
Moar, W. J., 1493
Moharramipour, S., 292, 1420, 2223
Moltó, E., 386
Montez, G. H., 1400
Moog, D. J., 1896
Moon, R. D., 1039
Moore, A., 1135
Morin, S., 2174
Morrell, J. J., 101
Morrow, J. L., 1071
Morse, J. G., 1172
Mosley, B. J., 1187
Mossadegh, M. S., 2112
Mottet, C., 1803
Muñoz, D., 17
Muegge, M. A., 735
Mueller, E. E., 1670
Muenworn, V., 1012
Mukawa, S., 257
Mullens, B. A., 550, 1929
Mullin, C., 1517
Muniruzzaman, S., 1647
Munyaneza, J. E., 1060
Murphy, A. F., 147
Murugan, M., 938
Mwanga, R.O.M., 1493
Mwatawala, M., 70
Mziray, H. A., 70

N

Nagle, H. T., 1636
Nagoshi, R. N., 360
Nakayama, S., 1564
Nansen, C., 219
Nault, B. A., 925, 1315
Navarro-Silva, M. A., 166
Naylor, R. A., 136
Neese, P., 603
Nelson, L. J., 770
Nelson, S. D., 1711
Neumann, G., 908
Neumann, P., 1, 10
Neven, L. G., 265
Newhouse, K. F., 119
Nguyen, B. K., 1517
Ni, X., 2072
Nichols, R. L., 158, 2174
Nielsen, D. G., 1187
Nishimoto, J., 2042
Nishisue, K., 1784
Nissinen, A., 1060
Noma, T., 583
Nozal, M.J.d., 1964
Nunez, S. C., 1937

O

Ode, P. J., 648
Odongo, B., 1493
Oh, H.-W., 25
Olsen, J., 416
Olson, D. M., 2072
O'Neil, R. J., 186
Onstad, D. W., 77, 849
Opit, G. P., 985, 1920
Orozco, G. V., 938
Orozco-Dávila, D., 85
Osbrink, W.L.A., 799
Otis, G., 708
Ottea, J., 869
Ouyang, Y., 1400

P

Paine, T. D., 1943
Palrang, D., 1711
Palyvos, N. E., 190
Pang, Y., 1832
Parajulee, M. N., 2009
Park, D.-S., 25
Park, H.-H., 563
Park, J.-J., 1474
Parker, A. G., 85
Parker, T. L., 472, 2147
Parman, V., 791
Pasqualini, E., 816
Pathan, A. K., 823
Peairs, F. B., 302, 1875
Pedrini, N., 1094
Pelz-Stelinski, K. S., 1531, 1972
Peng, Y.-F., 1444
Pereira, R., 85
Pereira, R. M., 1636
Pereira, Rui, 40
Pereira, S., 619
Perera, O. P., 861
Peretto, R., 816
Perring, T. M., 1474
Perumal, A., 1454
Pesek, J., 331
Peterson, C. J., 433
Peterson, R.K.D., 516
Petrice, T. R., 1682
Pettis, J. S., 10, 1517
Phillips, A. K., 448, 781
Piñero, J. C., 409, 1594
Picanço, M. C., 1438
Pierson, L. M., 1405
Pina, T., 386
Pineda, O., 2197

Pineda, S., 662
Pittendrigh, B. R., 203
Polk, D. F., 284
Poncet, C., 2242
Poppen, R. V., 1720
Potter, D. A., 1555
Pound, J. M., 211
Powell, Jr., R., 2072
Powis, K. J., 1347
Preckel, P. V., 1896
Progar, R. A., 242
Pryor, M. P., 1051
Puterka, G. J., 394, 958, 1875

Q

Qin, X.-W., 1665
Qin, Y., 2028
Qiu, B.-L., 1848
Quarcoo, F. Y., 762
Quinto, V., 2164

R

Raffa, K. F., 718
Ragsdale, D. W., 1483
Rahman, M. M., 1347
Rahman, T., 1744
Rajotte, E. G., 249
Ramírez-Lepe, M., 570
Ramírez-Suero, M., 570
Ramires, E. N., 166
Ramoutar, D., 1810
Randolph, T., 1875
Raphael, K. A., 1071
Raptopoulos, D., 1621
Rawi, C.S.M., 631
Razmjou, J., 966, 2223
Razzaq, M., 823
Reagan, T. E., 1431
Reagel, P. F., 641
Reay-Jones, F.P.F., 525
Rebek, E. J., 119
Reese, J. C., 938, 1405
Reeves, R. B., 525
Reineke, A., 898
Ren, S.-X., 1848
Requintina, Jr., E., 1810
Rhainds, M., 186
Rhea, R., 368
Rice, M. E., 54
Rice, S. J., 2155
Richmond, D. S., 1464
Rinderer, T. E., 917
Rinehart, J. P., 234
Rinella, M. J., 242
Ritter, W., 10
Rivi, M., 816
Roberts, P., 2236
Robertson, J. L., 1815
Rochon, K., 1019
Rodríguez, M. A., 482
Rodriguez, D., 1711
Rodriguez-Saona, C. R., 284
Rogers, D. J., 1206
Rogers, M. E., 1531
Rondon, S. I., 1306
Roubos, C. R., 1258
Roush, R. T., 1347
Royalty, R. N., 119, 726
Rudd, J. C., 1454
Rugman-Jones, P. F., 877, 1172
Rull, J., 2000
Rush, C. M., 1524
Rust, M. K., 808, 1761

S

Sétamou, M., 1711
Saad, A. R., 1770
Sabula, A., 597
Sadeghi, S. E., 2223
Sadof, C. S., 186
Saegerman, C., 1517
Sakamoto, H., 1784
Saldana, R., 1711
Salom, S. M., 1222
Saltzmann, K., 172
Sanderson, J. P., 1149
Sanon, A., 203
Sauphanor, B., 482
Sawyer, A. J., 1235
Sayyed, A. H., 823
Schäfer, M. O., 10
Schaffer, A., 228
Schal, C., 1636
Scharf, M. E., 1937
Scheffrahn, R. H., 1338
Schemerhorn, B. J., 2229
Schlenz, M., 1031
Schlyter, F., 1621
Schmidt, O., 1347
Schneider, M.-I., 662
Schotzko, D. J., 691
Schultheis, J. R., 2087
Schuphan, I., 1803
Schwarzlose, G., 1711
Scott, S. J., 1400
Scully, B. T., 2072
Segura, A., 619
Seiter, N. J., 1464
Seko, T., 1564
Selinger, L. B., 1019
Sengoda, V. G., 1060
Sensenbach, E. J., 1149
Shafir, S., 228
Shakya, S., 1086
Shatters, Jr., R. G., 2214
Shearman, D.C.A., 1071
Shelly, T., 2042
Shelton, A. M., 925, 1315, 2107
Shen, A., 843
Shen, Z., 2169
Shen, Z.-C., 1444
Sheppard, W. S., 917
Shi, G. L., 1372
Shipp, L., 563
Shiromoto, K., 1144, 2034
Shoemaker, R., 2042
Showler, A. T., 1180, 1431
Shukle, R. H., 2229
Shweil, S., 1621
Sial, A. A., 140, 340, 1277, 1378
Siegel, J. P., 1250
Siegfried, B. D., 1386
Silva, A.C.S., 166
Silva, J.J.D., 869
Silva-Zacarin, E.C.M.d., 676
Silverman, J., 1790
Sims, S. R., 448, 2118
Sisterson, M. S., 1051, 1821
Sivinski, J., 40
Smirle, M. J., 374
Smith, C. M., 938, 958
Smitley, D. R., 119
Soengas, P., 1866
Soleimannejad, S., 1420
Solera, M., 1493
Song, B. K., 1994
Sosa-Gomez, D. R., 869
Sotelo, G., 1860
Souder, S. K., 409
Soufbaf, M., 2019
Spafford, H., 1744
Spencer, T. A., 1386
Speybroeck, N., 1517
Spiewok, S., 1
Spivak, M., 1039
Sprick, P., 896
Stark, J. D., 409, 1594
Starkey, S., 938
Stary, P., 958
Steffey, K. L., 186
Steiner, H.-Y., 2187
Stelinski, L. L., 1972, 2100
Stock, S. P., 1821
Stodola, T. J., 1803
Stoltman, L., 1594
Storer, N. P., 1031
Stouthamer, R., 877, 1172
Su, N.-Y., 1327
Su, Z., 1832
Subramanyam, B., 1508
Suh, S.-J., 25
Suiter, D. R., 754
Sukonthabhirom, S., 1012
Sumerford, D. V., 249
Sumida, S., 676
Sunamura, E., 1784
Surridge, A.K.J., 887
Suzuki, S., 1784
Swanevelder, Z. H., 887
Sylvester, H. A., 917

T

Tabashnik, B. E., 1821
Tabuchi, R. L., 770
Tainchum, K., 1012
Takatsuki, J.-I., 1564

Tan, S. Y., 1386
Tanaka, Y., 1784
Tarpy, D. R., 1517
Tatsuki, S., 1784
Tawfik, H., 1621
Teal, P.E.A., 40
Tedders, W. L., 1
Tee, H.-S., 1770
Temple, J., 869
Terán-Vargas, A. P., 861
Terayama, M., 1784
Thomas, D. B., 1129
Thompson, D., 708
Thompson, G. D., 1031
Thompson, N., 1547
Thoms, E., 603
Throne, J. E., 197, 985, 1503
Tian, J.-C., 1444
Tillman, P. G., 2072
Tiwari, S., 1972, 2080
Tobin, P. C., 1569
Toews, M. D., 525, 991, 1002
Tomanović, S., 190
Tong, L., 1832
Tong, S., 492
Tuan, D. H., 127
Tuell, J. K., 557, 668

U

Ugine, T. A., 1149
Ulyshen, M. D., 1080
Urbaneja, A., 386

V

Vázquez, J. C., 726
Valle-Mora, J., 2000
Vandervoort, C., 1196
vanEngelsdorp, D., 1517
Vargas, R. I., 409, 1594
Vargo, E. L., 770, 791
Vasquez, E., 718
Vassiliou, V. A., 47
Vaughn, K., 219
Veillon, L., 1647
Velasco, P., 1866
Venter, E., 887
Vetter, R. S., 808, 1761
Villaverde, M. L., 1094
Vincent, C., 63
Viswanathan, P., 938
Voegtlin, D. J., 186, 509

W

Wagner, M. R., 1693
Walder, J.M.M., 85
Walker, G. P., 1172
Wallace, B. A., 2126
Walton, V. M., 416
Wang, B.-J., 2204
Wang, C., 172
Wang, H., 95
Wang, H. X., 1372
Wang, J.-J., 2204
Wang, M., 158
Wang, P., 492
Wang, Q.-X., 1166
Wang, X., 843
Wang, Y. N., 1372
Wang, Z.-T., 1365
War, M., 2042
Weaver, D. K., 516
Wei, D.-D., 2204
Weintraub, P. G., 1086
Weinzierl, R. A., 1815
Weisz, R., 2229
Weng, Y., 1454
West, M., 234
Westlake, T., 2140
White, W. H., 1431
Wiedenhoeft, A. C., 1682
Williams, T., 17, 577
Williamson, R. C., 1979
Willis, R. B., 1268, 2087
Wilson, J. P., 2072
Wiltz, B. A., 754
Winborne, I. C., 1950
Wise, J. C., 1196, 1720
Wistrom, C., 1051
Woloshuk, C., 1896
Wong, N., 437, 1754
Workneh, F., 1524
Wraight, S. P., 1149
Wratten, S. D., 2061
Wright, S. E., 394
Wu, J., 1100
Wu, K., 2169
Wu, Y., 843
Wu, J.-C., 1166

X

Xiong, M.-H., 1365
Xu, J., 1166

Y

Yali, M. P., 2223
Yang, W., 277
Yang, X.-B., 1729
Yaninek, S., 186
Ye, G.-Y., 1444
Ye, H., 95
Yee, W. L., 379
Yen, A. L., 2061
Yeselson, Y., 228
Yigezu, Y. A., 1896
Yocum, G. D., 234
Yoo, H.J.S., 186
Young, S., 835
Youngman, R. R., 2080
Younis, A. M., 2132
Yu, D., 277
Yuan, F.-H., 1665
Yuan, M.-L., 2204
Yuejin, W., 1988

Z

Zaborac, M., 2174
Zahiri, B., 292
Zalucki, M. P., 292, 1420, 2019
Zaman, F. U., 249
Zettler, J. L., 1950
Zhang, A., 284
Zhang, G., 277
Zhang, J., 1628, 1665
Zhang, K., 2204
Zhang, L., 492
Zhang, M., 1832
Zhang, R., 1100
Zhang, R.-J., 1628, 1665
Zhang, Y.-M., 1729
Zhang, Z., 95
Zhao, L. L., 1372
Zhou, Q., 1628
Zhu, L., 178
Zuñiga, E., 1860
Zurowski, C. L., 374

CUMULATIVE KEY WORD INDEX
Journal of Economic Entomology
Volume 103, 2010

A

abamectin,
  cyromazine, 2197
  Liriomyza trifolii, 2197
  random amplified polymorphic DNA polymerase chain reaction, 2197
  resistance, 2197
abamectin susceptibility,
  bioassays, 816
  insecticide resistance, 816
  pear psylla, 816
  pest control, 816
abandoned citrus,
  Asian citrus psyllid, 1972
  Candidatus Liberibacter asiaticus, 1972
  citrus greening, 1972
  huanglongbing, 1972
Abgrallaspis aguacatae,
  Aspidiotus nerii, 1172
  Diaspidiotus perniciosus, 1172
  Hemiberlesia lanatiae, 1172
  phoresy, 1172
acaricide resistance, 1355, 1360
  bioassay, 1360
  dieldrin, 1355
  GABA receptor, 1355
  PBO, 1360
  Rhipicephalus microplus, 1355
  synergist, 1360
  verbutin, 1360
acetamiprid, 379, 1815, 2107
  chlorantraniliprole, 1815
  Contarinia nasturtii, 2107
  egg laying, 379
  Grapholita molesta, 1815
  insecticide residue, 2107
  invasive species, 2107
  Rhagoletis indifferens, 379
  spinetoram, 1815
  spinosad, 1815
  spinosad bait, 379
  spray coverage, 2107
  thiamethoxam, 379
acoustic detection,
  eradication, 1135
  invasive species, 1135
acquisition,
  citrus greening disease, 1531
  huanglongbing, 1531
  inoculation, 1531
  vector, 1531
Acrobasis nuxvorella,
  biofix, 735
  Carya, 735
  heat unit, 735
  physiological model, 735
Acrobasis vacinii,
  life-stage activity, 1720
  parasitoid, 1720
  reduced-risk insecticide, 1720
action threshold,
  Allium cepa, 1315
  insecticide, 1315
  Thrips tabaci, 1315
Aculops lycopersici,
  Amblyseius swirskii, 563
  biological control, 563
  greenhouse tomato, 563
adult damage,
  Brachiaria, 1860
  host plant resistance, 1860
  life stages, 1860
  spittlebugs, 1860
adult emergence,
  density-dependent survival, 77
  Diabrotica virgifera virgifera, 77
  dose, 77
adults,
  insecticide resistance susceptibility, 2164
  larvae, 2164
  Thysanoptera, 2164
adventive range,
  Eucalyptus, 1943
  insect herbivores, 1943
  intentional introductions, 1943
  invasive species, 1943
Aethina tumida, 1, 10
  Apis mellifera, 1, 10
  entomopathogenic nematodes, 1
  Heterorhabditis spp., 1
  honey bee, 10
  Nosema ceranae, 10
  Steinernema spp., 1
  Varroa destructor, 10
Africa,
  Bactrocera latifrons, 70
  host preference, 70
  host range, 70
  Solanaceae, 70
aggregation,
  flight pattern, 95
  Pinus yunnanensis, 95
  Tomicus yunnanensis, 95
aggregation pheromone, 1603, 1915
  4-8-dimethyldecanal, 1915
  methyl (2*E*6*E*)-farnesoate, 1603
  methyl (*E*)-5-2610-trimethyl-59-undecadienoate, 1603
  methyl (*E*)-6-23-dihydrofarnesoate, 1603
  methyl geranate, 1603
  SPME, 1915
  stored products, 1915
aggression,
  Anoplolepis gracilipes, 1775
  interspecific, 1775
  intraspecific, 1775
  invasive species, 1775
agonism,
  interspecific, 1754
  intraspecific, 1754
  Microcerotermes crassus, 1754
  survival, 1754
Agrilus planipennis,
  borers, 119
  Fraxinus, 119
  imidacloprid, 119
Agrilus,
  Fraxinus, 603
  fumigation, 603
  quarantine, 603
  sulfuryl fluoride, 603
Agrotis ipsilon,
  Agrotis ipsilon nucleopolyhedrovirus, 1555
  baculovirus, 1555
  microbial control, 1555
  natural enemies, 1555
Agrotis ipsilon nucleopolyhedrovirus,
  Agrotis ipsilon, 1555
  baculovirus, 1555
  microbial control, 1555
  natural enemies, 1555
alfalfa weevil,
  fecundity, 292
  life table entropy, 292
  modeling, 292
  survivorship, 292
Allium cepa, 925, 1315
  action threshold, 1315
  insecticide, 1315
  Iris yellow spot virus, 925
  onion resistance, 925
  Thrips tabaci, 925, 1315
almonds,
  degree-days, 1250
  development, 1250
  navel orangeworm, 1250
  survival, 1250
Alpha radiation,
  Anastrepha fraterculus, 85
  Ceratitis capitata, 85
  irradiation, 85
  X radiation, 85
Alphitobius diaperinus,
  Australia, 2155
  darkling beetle, 2155
  pyrethroid cross-resistance, 2155
  topical application, 2155
alternate host,
  host suitability, 2009
  life table, 2009
  Lygus hesperus, 2009
  western tarnished plant bug, 2009
alternate host plants,
  citrus, 1051
  grape, 1051
  Pierce's disease, 1051
  vector control, 1051
alternative fumigation,
  Bambusa, 277
  Chlorophorus annularis, 277
  sulfuryl fluoride, 277

alternative pollinator,
Apis mellifera, 557
Bombus impatiens, 557
climate, 557
Vaccinium corymbosum, 557
aluminum-carboxymethylcellulose,
Bti, 570
México, 570
microbial control, 570
microencapsulation, 570
Amblyseius swirskii,
Aculops lycopersici, 563
biological control, 563
greenhouse tomato, 563
Amegilla,
flavor, 1286
horticulture, 1286
pollination, 1286
seeds, 1286
amplified fragment length polymorphism,
biogeography, 958
biotypes, 958
Diuraphis noxia, 958
isolation-by-distance, 958
Amyelois transitella,
Pyralidae, 314
Pyralis farinalis, 314
sex pheromone, 314
Anastrepha fraterculus,
Alpha radiation, 85
Ceratitis capitata, 85
irradiation, 85
X radiation, 85
Anastrepha ludens, 34, 1129, 2000
cold treatment, 2000
commodity treatment, 34, 1129
controlled atmosphere, 34
disinfestation, 34, 1129
Hass avocado, 2000
quarantine, 34, 1129
radiation, 1129
risk analysis, 2000
Tephritidae, 2000
Anastrepha suspensa,
Florida, 2214
host race formation, 2214
invasive insects, 2214
migration patterns, 2214
Anobiidae,
cold, 1909
heat, 1909
supercooling point, 1909
tolerance, 1909
Anomala orientalis,
Bacillaceae, 1547
Bacillus thuringiensis japonensis, 1547
Cyclocephala borealis, 1547
Scarabaeidae, 1547
Anopheles spp.,
collection methods, 973
French Guiana, 973
Anoplolepis gracilipes,
aggression, 1775
interspecific, 1775
intraspecific, 1775
invasive species, 1775
ant bait,
Malpighian tubules, 676
microscopy, 676
midgut, 676
postpharyngeal gland, 676
Anthonomus grandis grandis,
boll weevil, 356
eradication, 356
grandlure, 356
trapping, 356
antibiosis,
Aphis glycines, 1405
Glycine max, 1405
host plant resistance, 1405
tolerance, 1405
antifeedant,
oviposition deterrent, 1196
plant–insect–chemistry triad, 1196
sublethal effects, 1196
antifungal,
cellular encapsulation, 1327
epizootic, 1327
grooming, 1327
Isoptera, 1327
antiobiosis,
caffeic acid, 1438
caffeine, 1438
chlorogenic acid, 1438
Coffea, 1438
aphid, 374, 509
aphid resistance, 509
Aphis pomi, 374
Aphis spiraecola, 374
biotype, 509
insecticide resistance, 374
soybean, 509
soybean aphid, 509
susceptibility, 374
aphid resistance,
aphid, 509
biotype, 509
soybean, 509
soybean aphid, 509
Aphis glycines, 186, 648, 1405, 1483, 1670
antibiosis, 1405
Dowling, 648
electrical penetration graph, 648
field experiments, 1483
Glycine max, 1405
host plant resistance, 648, 1405, 1483
migration, 186
nonpersistent virus, 1670
Phaseolus vulgaris, 1670
predation, 1483
sampling, 1483
sampling procedure, 186
seasonal phenology, 1670
soybean aphid, 186, 648
suction trap, 186
tolerance, 1405
Aphis pomi,
aphids, 374
Aphis spiraecola, 374
insecticide resistance, 374
susceptibility, 374
Aphis spiraecola,
aphids, 374
Aphis pomi, 374
insecticide resistance, 374
susceptibility, 374
Aphthona,
biological control, 242
inundative release, 242
leafy spurge, 242
riparian area, 242
Apis mellifera, 1, 10, 228, 557, 1517, 1797, 1964
Aethina tumida, 1, 10
alternative pollinator, 557
Bombus impatiens, 557
classification and regression tree analysis, 1517
climate, 557
colony collapse disorder, 1517
colony losses, 1964
entomopathogenic nematodes, 1
epidemiology, 1517
Heterorhabditis spp., 1
honey bee, 10
monitoring insecticide resistance, 1797
Nosema ceranae, 10
pathogens, 1517
Persea americana, 228
perseitol, 228
pesticide residues, 1964
pollination, 228
Steinernema spp., 1
stored pollen, 1964
Vaccinium corymbosum, 557
Varroa destructor, 10, 1797
Apoidea,
insecticides, 668
nontarget effects, 668
Vaccinium corymbosum, 668
wild bee conservation, 668
apple,
conservation biological control, 1657
extrafloral nectar, 1657
Goniozus floridanus, 1657
Platynota idaeusalis, 1657
apple maggot,
behavior, 394
kaolin, 394
particle film technology, 394
Surround WP, 394
Aprostocetus hagenowii,
freeze, 1770
heat, 1770
parasitoid, 1770
Periplaneta americana, 1770
Arachnida,
brown recluse spider, 808
Loxosceles, 808
sericophily, 808
urban entomology, 808
Argentine ant,
control, 1784
Linepithema humile, 1784
synthetic trail pheromone, 1784
(Z)-9-hexadecenal, 1784
armored scales,
bark absorption, 1735
Christmas tree, 1735
IPM, 1735
systemic insecticide, 1735

artificial diet,
growth and reproduction, 17
OB yield, 17
SeMNPV, 17
Spodoptera exigua, 17
artificial infestation,
Brassica, 1866
glucosinolates, 1866
Mamestra brassicae, 1866
plant defense, 1866
artificial selection,
biological control, 1564
flightless, 1564
lady beetle, 1564
locomotor activity, 1564
Asian citrus psyllid,
abandoned citrus, 1972
Candidatus Liberibacter asiaticus, 1972
citrus greening, 1972
huanglongbing, 1972
Asian longhorned beetle,
biological invasions, 1569
emerald ash borer, 1569
firewood, 1569
invasive species management, 1569
Aspidiotus nerii,
Abgrallaspis aguacatae, 1172
Diaspidiotus perniciosus, 1172
Hemiberlesia lanatiae, 1172
phoresy, 1172
attractants,
ceralure, 1586
detection, 1586
Mediterranean fruit fly, 1586
trimedlure, 1586
attractive range,
codling moth, 1242
mark–release–recapture, 1242
mating disruption, 1242
spatial scale, 1242
Australia,
Alphitobius diaperinus, 2155
darkling beetle, 2155
pyrethroid cross-resistance, 2155
topical application, 2155
autoconfusion,
codlemone, 2100
Entostat powder, 2100
flight tunnel, 2100
mating disruption, 2100
avidin,
biotin, 497
pest control, 497
streptavidin, 497
transgenic plant, 497
avocado,
phytosanitary treatment, 1112
systems approach, 1112
Thaumatotibia leucotreta, 1112
azadirachtin,
emerald ash borer, 708
foliar uptake, 708
larval mortality, 708
systemic injection, 708

B

B biotype,
Bemisia tabaci, 158
cytochrome P450, 158
glutathione S-transferase, 158
monooxygenase, 158
resistance to pyriproxyfen, 158
B. carambolae,
Bactrocera payayae, 1994
cytochrome oxidase I, 1994
single-nucleotide polymorphism-polymerase chain reaction, 1994
species differentiation, 1994
Bacillaceae,
Anomala orientalis, 1547
Bacillus thuringiensis japonensis, 1547
Cyclocephala borealis, 1547
Scarabaeidae, 1547
Bacillus thuringiensis japonensis,
Anomala orientalis, 1547
Bacillaceae, 1547
Cyclocephala borealis, 1547
Scarabaeidae, 1547
Bacillus thuringiensis, 1019, 1444, 2147, 2187
Bt, 2187
Chilo suppressalis, 1444
Cry1Ab/Vip3H protein, 1444
Cry1Ac, 2147
Cry2Ab, 2147
Diabrotica virgifera virgifera, 2187
Helicoverpa punctigera, 2147
horn fly, 1019
mCry3A, 2187
microbial control, 1019
resistance, 2147
resistance management, 2187
Sesamia inferens, 1444
stable fly, 1019
transgenic rice, 1444
bacterial diversity,
colonization, 1832
ethidium monoazide, 1832
Musca domestica, 1832
16S rDNA, 1832
Bactericera cockerelli,
life table, 1729
mortality, 1729
population dynamics, 1729
Zebra Chip, 1729
Bactrocera cucurbitae, 409, 1594
Bactrocera dorsalis, 409
control, 1594
male annihilation treatment, 1594
Min-U-Gel, 1594
naled, 409, 1594
spinosad, 409
tephritid flies, 409
Bactrocera dorsalis,
Bactrocera cucurbitae, 409
naled, 409
spinosad, 409
tephritid flies, 409
Bactrocera jarvisi,
Bactrocera tryoni, 1071
genetic markers, 1071
interspecific hybridization, 1071
Bactrocera latifrons,
Africa, 70
host preference, 70
host range, 70
Solanaceae, 70
Bactrocera payayae,
B. carambolae, 1994
cytochrome oxidase I, 1994
single-nucleotide polymorphism-polymerase chain reaction, 1994
species differentiation, 1994
Bactrocera tryoni,
Bactrocera jarvisi, 1071
genetic markers, 1071
interspecific hybridization, 1071
baculovirus,
Agrotis ipsilon, 1555
Agrotis ipsilon nucleopolyhedrovirus, 1555
microbial control, 1555
natural enemies, 1555
bagging,
chemical control, 1577
sanitation, 1577
systems approach, 1577
baiting,
chitin synthesis inhibitor, 423
Coptotermes, 423
nest temperature, 423
peritrophic membrane, 423
bamboo,
cost/benefit, 101
Dinoderus minutus, 101
prophylactic treatment, 101
wood preservatives, 101
Bambusa,
alternative fumigation, 277
Chlorophorus annularis, 277
sulfuryl fluoride, 277
barcoding,
cytochrome c oxidase I gene, 877
molecular identification, 877
28SD2, 877
tospovirus vector, 877
bark absorption,
armored scales, 1735
Christmas tree, 1735
IPM, 1735
systemic insecticide, 1735
bark beetles,
exotic species, 1187
Ohio, 1187
phenology, 1187
semiochemicals, 1187
barley,
biotypes, 938
Diuraphis noxia, 938
resistance, 938
RWA, 938
barrier,
horizontal toxicity, 754
insecticide, 754
Solenopsis invicta, 754
baseline susceptibility,
chlorantraniliprole, 843
discriminating dose, 843
Plutella xylostella, 843
resistance monitoring, 843

Beauveria bassiana,
diet, 1542
starvation, 1542
stress, 1542
Tribolium castaneum, 1542
bed bug, 136, 172, 1636
Cimex lectularius, 136, 172
cockroach, 1636
control, 136
dispersal, 172
drugstore beetle, 1636
laundry, 136
monitoring, 172
red flour beetle, 1636
rice weevil, 1636
behavior,
apple maggot, 394
kaolin, 394
particle film technology, 394
Surround WP, 394
Bemisia tabaci, 158, 1848, 2174
B biotype, 158
cytochrome P450, 158
glutathione S-transferase, 158
infection frequency, 1848
insect growth regulators, 2174
insecticide resistance, 2174
monooxygenase, 158
neonicotinoids, 2174
parasitoid, 1848
predator, 1848
Q biotype, 2174
resistance to pyriproxyfen, 158
Wolbachia supergroups, 1848
benzaldehyde,
Circle trap, 744
grandisoic acid, 744
plum curculio, 744
pyramid trap, 744
benzoyl urea insect growth regulator,
chitin synthesis inhibitor, 770
subterranean termites, 770
termite baits, 770
termite colony elimination, 770
binomial sampling plan,
Ectomyelois ceratoniae, 1474
Phoenix dactylifera, 1474
resampling validation, 1474
seasonal population dynamics, 1474
bioassay, 482, 1360
acaricide resistance, 1360
Cydia pomonella, 482
detoxifying enzymes, 482
organophosphates, 482
PBO, 1360
resistance detection, 482
synergist, 1360
verbutin, 1360
bioassays,
abamectin susceptibility, 816
insecticide resistance, 816
pear psylla, 816
pest control, 816
biocontrol, 583, 908, 1149
Bradysia, 1149
Coenosia attenuata, 1149
cytochrome oxidase subunit I gene, 908
development, 1149
DNA fingerprinting, 908
endoparasitoids, 908
molecular markers, 908
parasitoids, 583
predator, 1149
predatory flies, 583
soybean aphid, 583
biofix,
Acrobasis nuxvorella, 735
Carya, 735
heat unit, 735
physiological model, 735
biogeography,
amplified fragment length polymorphism, 958
biotypes, 958
Diuraphis noxia, 958
isolation-by-distance, 958
biological control, 242, 308, 416, 563, 1094, 1564, 2061
Aculops lycopersici, 563
Amblyseius swirskii, 563
Aphthona, 242
artificial selection, 1564
Calepitrimerus vitis, 2061
Colomerus vitis, 2061
defensive pheromone, 1094
entomopathogenic fungi, 1094
entomopathogenic nematodes, 416
flightless, 1564
greenhouse tomato, 563
hazelnut, 416
insecticide, 308
integrated pest management in viticulture, 2061
inundative release, 242
lady beetle, 1564
leafy spurge, 242
locomotor activity, 1564
nematode efficacy, 416
orchard floor cover, 416
Phytoseiidae, 2061
riparian area, 242
solid phase microextraction, 1094
stored products pests, 1094
sublethal effects, 308
Trichogramma evanescens, 308
Trichogramma semblidis, 308
biological invasions,
Asian longhorned beetle, 1569
emerald ash borer, 1569
firewood, 1569
invasive species management, 1569
biomass production,
Lymantria dispar, 718
microbial pesticides, 718
Populus, 718
short rotation trees, 718
biopesticide,
Callosobruchus maculatus, 203
cowpea storage, 203
spinosad, 203
biotic,
herbivory, 516
phosphorus deficiency, 516
stem mining, 516
water stress, 516
biotin,
avidin, 497
pest control, 497
streptavidin, 497
transgenic plant, 497
biotype, 509, 887, 938, 949, 958, 1454
aphid, 509
aphid resistance, 509
amplified fragment length polymorphism, 958
barley, 938
biogeography, 958
detached leaves, 949
Diuraphis noxia, 887, 938, 958
endosymbiont variation, 887
genetic diversity, 1454
isolation-by-distance, 958
microsatellite, 1454
resistance, 938
RWA, 938
Schizaphis graminum, 1454
simple sequence repeat, 1454
soybean, 509
soybean aphid, 509, 949
soybean host plant resistance, 949
Blattella germanica, 448, 460, 781, 2118
essential oils, 448, 781
formulation, 2118
fumigation, 781
insecticide resistance, 460
microencapsulation, 2118
ootheca, 448, 781
repellency, 2118
resistance mechanism, 460
Singapore, 460
synergism, 460
toxicity, 448, 2118
Blissus insularis,
chinch bug, 726
insect diet, 726
rearing methods, 726
blueberries,
Dasineura oxycoccana, 1258
distribution, 1258
emergence trap, 1258
rabbiteye, 1258
boll damage,
cotton, 2236
external lesions, 2236
sampling, 2236
stink bug, 2236
boll weevil,
Anthonomus grandis grandis, 356
eradication, 356
grandlure, 356
trapping, 356
Bollgard II,
chilling day-degrees, 1206
critical period, 1206
seasonal production, 1206
temperature threshold, 1206
Bombus impatiens,
alternative pollinator, 557
Apis mellifera, 557
climate, 557
Vaccinium corymbosum, 557
Boophilus annulatus,
Boophilus microplus, 211
cattle fever tick, 211

Rhipicephalus, 211
white-tailed deer, 211
Boophilus microplus,
Boophilus annulatus, 211
cattle fever tick, 211
Rhipicephalus, 211
white-tailed deer, 211
borers,
Agrilus planipennis, 119
Fraxinus, 119
imidacloprid, 119
boring behavior,
Chilo suppressalis, 1412
plant resistance, 1412
rice, 1412
silicon, 1412
botanical acaricide,
essential oil, 1293
natural fumigant, 1293
Neoseiulus californicus, 1293
Tetranychus urticae, 1293
Brachiaria,
adult damage, 1860
host plant resistance, 1860
life stages, 1860
spittlebugs, 1860
Bradysia,
biocontrol, 1149
Coenosia attenuata, 1149
development, 1149
predator, 1149
Brassica,
artificial infestation, 1866
glucosinolates, 1866
Mamestra brassicae, 1866
plant defense, 1866
Brevicoryne brassicae,
camptothecin, 492
Chilo suppressalis, 492
natural insecticide, 492
Nilaparvata lugens, 492
BROTE,
downy brome, 348
Neoseiulus, 348
spider mites, 348
western corn rootworm, 348
brown recluse spider,
Arachnida, 808
Loxosceles, 808
sericophily, 808
urban entomology, 808
Bt,
Bacillus thuringiensis, 2187
Diabrotica virgifera virgifera, 2187
mCry3A, 2187
resistance management, 2187
Bt bioassays,
Bt toxins, 1493
Ipomoea batatas, 1493
Bt corn,
refuge-in-a-bag, 147
resistance management, 147
seed mix refuge, 147
western corn rootworm, 147
Bt maize, 1394, 1803
Cry1Ab, 1394
Cry3Bb, 1394
European corn borer, 1803
HDR strategy, 1803
insect resistance evolution, 1394
Mon810, 1803
resistance management, 1803
transgenic maize, 1394
Bt resistance, 472, 835
Cry2Ab, 835
F_2 and F_1 screens, 472
fitness costs, 835
Helicoverpa armigera, 472, 835
Bt toxins,
Bt bioassays, 1493
Ipomoea batatas, 1493
Bti,
aluminum-carboxymethylcellulose, 570
México, 570
microbial control, 570
microencapsulation, 570

C

caffeic acid,
antiobiosis, 1438
caffeine, 1438
chlorogenic acid, 1438
Coffea, 1438
caffeine,
antiobiosis, 1438
caffeic acid, 1438
chlorogenic acid, 1438
Coffea, 1438
Calepitrimerus vitis,
biological control, 2061
Colomerus vitis, 2061
integrated pest management in viticulture, 2061
Phytoseiidae, 2061
California least tern,
fipronil, 1761
horizontal transfer, 1761
Linepithema humile, 1761
Callosobruchus maculatus,
biopesticide, 203
cowpea storage, 203
spinosad, 203
camptothecin,
Brevicoryne brassicae, 492
Chilo suppressalis, 492
natural insecticide, 492
Nilaparvata lugens, 492
Candidatus Liberibacter asiaticus
abandoned citrus, 1972
Asian citrus psyllid, 1972
citrus greening, 1972
huanglongbing, 1972
cannibalism,
juvenile hormone, 577
nucleopolyhedrovirus production, 577
Spodoptera exigua, 577
canola,
IPM, 654
Lepidoptera, 654
pheromone monitoring, 654
Plutella xylostella, 654
canola cultivars,
life-table analysis, 2019
model based survival, 2019
Plutella xylostella, 2019
canopy distribution,
colored sticky cards, 47
grapefruit-lemon, 47
host plants, 47
Kelly's citrus thrips, 47
canopy malaise,
exotic species survey, 698
species richness, 698
trapping, 698
canopy translocation,
concentration, 368
eastern hemlock, 368
hemlock woolly adelgid, 368
imidacloprid, 368
carbohydrate,
Coptotermes formosanus, 1647
2-deoxy-d-galactose, 1647
sugar, 1647
termite, 1647
Caribbean fruit fly, 40, 533
hydrolyzed yeast, 40
male competitiveness, 40
methoprene, 40
sexual performance, 40
spatial analysis, 533
synthetic lures, 533
torula yeast, 533
trapping, 533
Carposina niponensis,
fumigant, 1988
phosphine, 1988
toxicity, 1988
carrot,
liberibacter, 1060
plant pathogen, 1060
psyllid, 1060
Trioza apicalis, 1060
Carya,
Acrobasis nuxvorella, 735
biofix, 735
heat unit, 735
physiological model, 735
cattle fever tick,
Boophilus annulatus, 211
Boophilus microplus, 211
Rhipicephalus, 211
white-tailed deer, 211
CATTS,
controlled atmosphere, 265
false codling moth, 265
heat, 265
postharvest, 265
cellular encapsulation,
antifungal, 1327
epizootic, 1327
grooming, 1327
Isoptera, 1327
ceralure,
attractants, 1586
detection, 1586
Mediterranean fruit fly, 1586
trimedlure, 1586
Cerambycidae,
Fluon, 641
pheromone, 641
trap, 641
wood-boring insect, 641
Ceratitis capitata, 85, 1100
Alpha radiation, 85
Anastrepha fraterculus, 85
Ceratitis cosyra, 1100
Ceratitis rosa, 1100
China, 1100

irradiation, 85
risk analysis, 1100
X radiation, 85
Ceratitis cosyra,
Ceratitis capitata, 1100
Ceratitis rosa, 1100
China, 1100
risk analysis, 1100
Ceratitis rosa,
Ceratitis capitata, 1100
Ceratitis cosyra, 1100
China, 1100
risk analysis, 1100
Ceratitis species,
Dacus species, 1842
mitochondrial DNA, 1842
molecular taxonomy, 1842
Chaitophorus leucomelas,
life table, 2223
Populus, 2223
susceptibility, 2223
chemical barrier,
fipronil, 433
imidacloprid, 433
soil penetration, 433
termites, 433
chemical control,
bagging, 1577
sanitation, 1577
systems approach, 1577
chilling day-degrees,
Bollgard II, 1206
critical period, 1206
seasonal production, 1206
temperature threshold, 1206
Chilo suppressalis, 492, 1412, 1444
Bacillus thuringiensis, 1444
boring behavior, 1412
Brevicoryne brassicae, 492
camptothecin, 492
cry1Ab/Vip3H protein, 1444
natural insecticide, 492
Nilaparvata lugens, 492
plant resistance, 1412
rice, 1412
Sesamia inferens, 1444
silicon, 1412
transgenic rice, 1444
China,
Ceratitis capitata, 1100
Ceratitis cosyra, 1100
Ceratitis rosa, 1100
risk analysis, 1100
chinch bug,
Blissus insularis, 726
insect diet, 726
rearing methods, 726
chitin synthesis inhibitor, 423, 770
baiting, 423
benzoyl urea insect growth regulator, 770
Coptotermes, 423
nest temperature, 423
peritrophic membrane, 423
subterranean termites, 770
termite baits, 770
termite colony elimination, 770
chlorantraniliprole, 140, 843, 1277, 1378, 1815
acetamiprid, 1815
baseline susceptibility, 843
Choristoneura rosaceana, 1277
discriminating dose, 843
exposure period, 1277
field-collected populations, 140
Grapholita molesta, 1815
heritability, 1378
insecticide resistance, 140
obliquebanded leafroller, 140, 1378
Plutella xylostella, 843
residues, 1277
resistance, 1378
resistance monitoring, 843
spinetoram, 140, 1277, 1378, 1815
spinosad, 1815
chlorogenic acid,
antiobiosis, 1438
caffeic acid, 1438
caffeine, 1438
Coffea, 1438
Chlorophorus annularis,
alternative fumigation, 277
Bambusa, 277
sulfuryl fluoride, 277
Choristoneura rosaceana,
chlorantraniliprole, 1277
exposure period, 1277
residues, 1277
spinetoram, 1277
Christmas tree,
armored scales, 1735
bark absorption, 1735
IPM, 1735
systemic insecticide, 1735
Chrsoperla carnea,
fitness benefit, 823
organophosphate, 823
pyrethroids, 823
resistance, 823
Cimex lectularius, 136, 172
bed bug, 136, 172
control, 136
dispersal, 172
laundry, 136
monitoring, 172
Circle trap,
benzaldehyde, 744
grandisoic acid, 744
plum curculio, 744
pyramid trap, 744
citrus, 1051, 1400
alternate host plants, 1051
grape, 1051
insecticide resistance, 1400
organophosphate, 1400
Pierce's disease, 1051
vector control, 1051
citrus greening, 127, 1972
abandoned citrus, 1972
Asian citrus psyllid, 1972
Candidatus Liberibacter asiaticus, 1972
huanglongbing, 127, 1972
king mandarin, 127
population, 127
systemic insecticide, 127
citrus greening disease, 541, 1531
acquisition, 1531
huanglongbing, 541, 1531
inoculation, 1531
monitoring, 541
vector, 1531
citrus mealybug,
grapevine, 619
Grapevine leafroll-associated virus 3, 619
insect, 619
Planococcus citri, 619
classical biological control,
concealed host, 1080
exotic, 1080
forest pest, 1080
non-native, 1080
classification and regression tree analysis,
Apis mellifera, 1517
colony collapse disorder, 1517
epidemiology, 1517
pathogens, 1517
climate,
alternative pollinator, 557
Apis mellifera, 557
Bombus impatiens, 557
Vaccinium corymbosum, 557
Cnaphalocrocis medinalis,
resistance, 1166
rice variety, 1166
Coccidae,
Linepithema humile, 1790
Mesolecanium nigrofasciatum, 1790
pest management, 1790
red maple, 1790
cockroach,
bed bug, 1636
drugstore beetle, 1636
red flour beetle, 1636
rice weevil, 1636
codlemone, 1704, 2100
autoconfusion, 2100
Cydia pomonella, 1704
Entostat powder, 2100
flight tunnel, 2100
mating disruption, 2100
methoxyfenozide, 1704
responsiveness, 1704
sublethal effects, 1704
codling moth,
attractive range, 1242
mark–release–recapture, 1242
mating disruption, 1242
spatial scale, 1242
Coenosia attenuata,
biocontrol, 1149
Bradysia, 1149
development, 1149
predator, 1149
Coffea,
antiobiosis, 1438
caffeic acid, 1438
caffeine, 1438
chlorogenic acid, 1438
Coffea arabica,
Hypothenemus hampei, 1159
infestation, 1159
reproduction, 1159
temperature, 1159

cold,
Anobiidae, 1909
heat, 1909
supercooling point, 1909
tolerance, 1909
cold treatment,
Anastrepha ludens, 2000
Hass avocado, 2000
risk analysis, 2000
Tephritidae, 2000
Coleoptera,
Cylas formicarius elegantulus, 1144
laboratory adaptation, 1144
sterile insect technique, 1144
collection methods,
Anopheles spp., 973
French Guiana, 973
Colomerus vitis,
biological control, 2061
Calepitrimerus vitis, 2061
integrated pest management in viticulture, 2061
Phytoseiidae, 2061
colonization,
bacterial diversity, 1832
ethidium monoazide, 1832
Musca domestica, 1832
16S rDNA, 1832
colony collapse disorder,
Apis mellifera, 1517
classification and regression tree analysis, 1517
epidemiology, 1517
pathogens, 1517
colony elimination,
colony suppression, 791
microsatellites, 791
Reticulitermes flavipes, 791
Reticulitermes virginicus, 791
colony losses,
Apis mellifera, 1964
pesticide residues, 1964
stored pollen, 1964
colony suppression,
colony elimination, 791
microsatellites, 791
Reticulitermes flavipes, 791
Reticulitermes virginicus, 791
colored sticky cards,
canopy distribution, 47
grapefruit-lemon, 47
host plants, 47
Kelly's citrus thrips, 47
commodity treatment, 34, 1129, 1950
Anastrepha ludens, 34, 1129
controlled atmosphere, 34
disinfestation, 34, 1129, 1950
irradiation, 1950
quarantine, 34, 1129, 1950
radiation, 1129
complementary log-log modeling,
pesticide, 1665
rice insect pest, 1665
virulence, 1665
concealed host,
classical biological control, 1080
exotic, 1080
forest pest, 1080
non-native, 1080
concentration,
canopy translocation, 368
eastern hemlock, 368
hemlock woolly adelgid, 368
imidacloprid, 368
Conoderus vespertinus,
sweetpotato, 1268
tobacco wireworm, 1268
wireworm, 1268
Conoderus spp.,
crop rotation, 2087
Melanotus communis, 2087
wireworm, 2087
conophthorin,
Gnathotrichus materiarius, 2094
Ips grandicollis, 2094
Ips pini, 2094
Xylosandrus germanus, 2094
conservation biological control,
apple, 1657
extrafloral nectar, 1657
Goniozus floridanus, 1657
Platynota idaeusalis, 1657
Contarinia nasturtii,
acetamiprid, 2107
insecticide residue, 2107
invasive species, 2107
spray coverage, 2107
contour analysis,
geostatistics, 1886
Mediterranean fruit fly, 1886
spatial statistics, 1886
variogram, 1886
control, 136, 1594, 1784
Argentine ant, 1784
Bactrocera cucurbitae, 1594
bed bug, 136
Cimex lectularius, 136
laundry, 136
Linepithema humile, 1784
male annihilation treatment, 1594
Min-U-Gel, 1594
naled, 1594
synthetic trail pheromone, 1784
(Z)-9-hexadecenal, 1784
control level,
Diaphorina citri, 1711
imidacloprid, 1711
Phyllocnistis citrella, 1711
systemic uptake, 1711
controlled atmosphere, 34, 265, 272
Anastrepha ludens, 34
CATTS, 265
commodity treatment, 34
disinfestation, 34
false codling moth, 265
grape benchgraft, 272
heat, 265
mealybug, 272
Planococcus ficus, 272
postharvest, 265
quarantine, 34
ultralow oxygen, 272
Coptotermes formosanus, 799, 1338, 1647, 2132
carbohydrate, 1647
Coptotermes gestroi, 1338
2-deoxy-d-galactose, 1647
desiccation, 799
food consumption, 2132
naphthalene derivatives, 2132
repellency, 2132
seaborne, 1338
spatial analysis, 1338
sugar, 1647
termite, 1647
termite infestation, 1338
tunneling, 799
wood consumption, 799
Coptotermes gestroi, 437, 1338
Coptotermes formosanus, 1338
Microcerotermes crassus, 437
moisture, 437
seaborne, 1338
spatial analysis, 1338
termite infestation, 1338
tunneling activity, 437
wood consumption, 437
Coptotermes,
baiting, 423
chitin synthesis inhibitor, 423
nest temperature, 423
peritrophic membrane, 423
cost/benefit,
bamboo, 101
Dinoderus minutus, 101
prophylactic treatment, 101
wood preservatives, 101
cotton,
boll damage, 2236
external lesions, 2236
sampling, 2236
stink bug, 2236
coverage,
integrated pest management, 386
mineral oil, 386
twospotted spider mite, 386
cowpea storage,
biopesticide, 203
Callosobruchus maculatus, 203
Spinosad, 203
critical period,
Bollgard II, 1206
chilling day-degrees, 1206
seasonal production, 1206
temperature threshold, 1206
crop rotation, 1464, 2087
Conoderus spp., 2087
image analysis, 1464
Melanotus communis, 2087
soybean herbivory, 1464
western corn rootworm, 1464
wireworm, 2087
cross-tolerance,
Cry1Ac, 2169
Helicoverpa armigera, 2169
Vip3Aa, 2169
Cry1Ab, 54, 1386, 1394
Bt maize, 1394
Cry3Bb, 1394
fitness costs, 1386
high-dose refuge, 1386
insect resistance evolution, 1394
intraguild competition, 54
MON810, 54
resistance and insect resistance management, 1386
Striacosta albicosta, 54
transgenic maize, 1394
YieldGard, 54

Cry1Ab/Vip3H protein,
Bacillus thuringiensis, 1444
Chilo suppressalis, 1444
Sesamia inferens, 1444
transgenic rice, 1444
Cry1Ac, 2147, 2169
Bacillus thuringiensis, 2147
cross-tolerance, 2169
Cry2Ab, 2147
Helicoverpa armigera, 2169
Helicoverpa punctigera, 2147
resistance, 2147
Vip3Aa, 2169
Cry1F,
fall armyworm, 1031
transgenic maize, 1031
Cry2Ab, 835, 2147
Bacillus thuringiensis, 2147
Bt resistance, 835
Cry1Ac, 2147
fitness costs, 835
Helicoverpa armigera, 835
Helicoverpa punctigera, 2147
resistance, 2147
Cry3Bb,
Bt maize, 1394
Cry1Ab, 1394
insect resistance evolution, 1394
transgenic maize, 1394
Curculio caryae,
new host association, 1312
quarantine threat, 1312
weevil DNA, 1312
Cyclocephala borealis,
Anomala orientalis, 1547
Bacillaceae, 1547
Bacillus thuringiensis japonensis, 1547
Scarabaeidae, 1547
Cydia pomonella, 482, 1704
bioassay, 482
codlemone, 1704
detoxifying enzymes, 482
methoxyfenozide, 1704
organophosphates, 482
resistance detection, 482
responsiveness, 1704
sublethal effects, 1704
Cylas formicarius elegantulus, 1144, 2034
Coleoptera, 1144
gamma radiation, 2034
laboratory adaptation, 1144
mating behavior, 2034
sterile insect technique (SIT), 1144, 2034
substerilization, 2034
Cylindrocopturus adspersus,
degree-day model, 302
predictive model, 302
sunflower stem weevil, 302
cyromazine,
abamectin, 2197
Liriomyza trifolii, 2197
random amplified polymorphic DNA polymerase chain reaction, 2197
resistance, 2197
cytochrome c oxidase I gene,
barcoding, 877
molecular identification, 877
28SD2, 877
tospovirus vector, 877
cytochrome oxidase I,
B. carambolae, 1994
Bactrocera payayae, 1994
single-nucleotide polymorphism-polymerase chain reaction, 1994
species differentiation, 1994
cytochrome oxidase subunit I gene,
biocontrol, 908
DNA fingerprinting, 908
endoparasitoids, 908
molecular markers, 908
cytochrome oxidase subunit II,
molecular diagnostic marker, 898
pest management, 898
weevil, 898
cytochrome P450,
B biotype, 158
Bemisia tabaci, 158
glutathione S-transferase, 158
monooxygenase, 158
resistance to pyriproxyfen, 158

D

Dacus species,
Ceratitis species, 1842
mitochondrial DNA, 1842
molecular taxonomy, 1842
dairy pest management,
insecticide bait, 1937
insecticide resistance, 1937
integrated pest management, 1937
Musca domestica, 1937
darkling beetle,
Alphitobius diaperinus, 2155
Australia, 2155
pyrethroid cross-resistance, 2155
topical application, 2155
Dasineura oxycoccana,
blueberries, 1258
distribution, 1258
emergence trap, 1258
rabbiteye, 1258
defensive pheromone,
biological control, 1094
entomopathogenic fungi, 1094
solid phase microextraction, 1094
stored products pests, 1094
degree-day model,
Cylindrocopturus adspersus, 302
predictive model, 302
sunflower stem weevil, 302
degree-days,
almonds, 1250
development, 1250
navel orangeworm, 1250
survival, 1250
demographic parameters,
Helicoverpa armigera, 1420
nutritional indices, 1420
soybean cultivars, 1420
demography,
population growth parameter, 966
potato cultivar, 966
potato tuberworm, 966
Solanum tuberosum, 966
Dendroctonus,
Ips pini, 1693
microclimate, 1693
Pinus ponderosa, 1693
thinning, 1693
density-dependent survival,
adult emergence, 77
Diabrotica virgifera virgifera, 77
dose, 77
2-deoxy-d-galactose,
carbohydrate, 1647
Coptotermes formosanus, 1647
sugar, 1647
termite, 1647
desiccation, 799, 1306
Coptotermes formosanus, 799
insecticide application timing, 1306
integrated pest management, 1306
irrigation, 1306
Phthorimaea operculella, 1306
tunneling, 799
wood consumption, 799
detached leaves,
biotypes, 949
soybean aphid, 949
soybean host plant resistance, 949
detection,
attractants, 1586
ceralure, 1586
Mediterranean fruit fly, 1586
trimedlure, 1586
detoxifying enzymes,
bioassay, 482
Cydia pomonella, 482
organophosphates, 482
resistance detection, 482
development, 662, 1149, 1250
almonds, 1250
biocontrol, 1149
Bradysia, 1149
Coenosia attenuata, 1149
degree-days, 1250
fecundity, 662
fertility, 662
methoxyfenozide, 662
navel orangeworm, 1250
predator, 1149
Spodoptera exigua, 662
survival, 1250
developmental period,
feeding, 1744
Frankliniella occidentalis, 1744
oviposition, 1744
survival rate, 1744
developmental rates,
population growth, 1920
stored grain, 1920
stored products, 1920
Diabrotica virgifera virgifera, 77, 2052, 2187
adult emergence, 77
Bacillus thuringiensis, 2187
Bt, 2187
density-dependent survival, 77
dose, 77
insect behavior, 2052
larvae, 2052
mCry3A, 2187

resistance, 2052
resistance management, 2187
Zea mays, 2052
Diaphorina citri,
control level, 1711
imidacloprid, 1711
Phyllocnistis citrella, 1711
systemic uptake, 1711
Diaspidiotus perniciosus,
Abgrallaspis aguacatae, 1172
Aspidiotus nerii, 1172
Hemiberlesia lanatiae, 1172
phoresy, 1172
diatomaceous earth,
storage mite pests, 190
stored wheat, 190
Tyrophagus putrescentiae, 190
dieldrin,
acaricide resistance, 1355
GABA receptor, 1355
Rhipicephalus microplus, 1355
diet,
Beauveria bassiana, 1542
starvation, 1542
stress, 1542
Tribolium castaneum, 1542
4-8-dimethyldecanal,
aggregation pheromone, 1915
SPME, 1915
stored products, 1915
Dinoderus minutus,
bamboo, 101
cost/benefit, 101
prophylactic treatment, 101
wood preservatives, 101
Dioryctria amatella,
Dioryctria clarioralis, 1216
Dioryctria disclusa, 1216
Dioryctria merkeli, 1216
(3Z6Z9Z12Z15Z)-pentacosapentaene, 1216
Dioryctria clarioralis,
Dioryctria amatella, 1216
Dioryctria disclusa, 1216
Dioryctria merkeli, 1216
(3Z6Z9Z12Z15Z)-pentacosapentaene, 1216
Dioryctria disclusa,
Dioryctria amatella, 1216
Dioryctria clarioralis, 1216
Dioryctria merkeli, 1216
(3Z6Z9Z12Z15Z)-pentacosapentaene, 1216
Dioryctria merkeli,
Dioryctria amatella, 1216
Dioryctria clarioralis, 1216
Dioryctria disclusa, 1216
(3Z6Z9Z12Z15Z)-pentacosapentaene, 1216
discriminating dose,
baseline susceptibility, 843
chlorantraniliprole, 843
Plutella xylostella, 843
resistance monitoring, 843
disinfestation, 34, 1129, 1950
Anastrepha ludens, 34, 1129
commodity treatment, 34, 1129, 1950
controlled atmosphere, 34
irradiation, 1950
quarantine, 34, 1129, 1950
radiation, 1129
dispersal, 63, 172, 331, 1524
bed bug, 172
Cimex lectularius, 172
monitoring, 172
movement, 63
Ostrinia nubilalis, 331
path, 63
plant abandonment, 331
plant disease vector, 1524
psyllid yellows, 1524
silking behavior, 331
tracking, 63
transgenic corn, 331
zebra chip, 1524
distribution,
blueberries, 1258
Dasineura oxycoccana, 1258
emergence trap, 1258
rabbiteye, 1258
Diuraphis frequens,
Diuraphis mexicana, 1875
Diuraphis noxia, 1875
Diuraphis tritici, 1875
invasive species, 1875
Diuraphis mexicana,
Diuraphis frequens, 1875
Diuraphis noxia, 1875
Diuraphis tritici, 1875
invasive species, 1875
Diuraphis noxia, 887, 938, 958, 1875
amplified fragment length polymorphism, 958
barley, 938
biogeography, 958
biotypes, 887, 938, 958
Diuraphis frequens, 1875
Diuraphis mexicana, 1875
Diuraphis tritici, 1875
endosymbiont variation, 887
invasive species, 1875
isolation-by-distance, 958
resistance, 938
RWA, 938
Diuraphis tritici,
Diuraphis frequens, 1875
Diuraphis mexicana, 1875
Diuraphis noxia, 1875
invasive species, 1875
DNA fingerprinting,
biocontrol, 908
cytochrome oxidase subunit I gene, 908
endoparasitoids, 908
molecular markers, 908
dose,
adult emergence, 77
density-dependent survival, 77
Diabrotica virgifera virgifera, 77
dose–mortality analysis,
dose–mortality analysis, 869
neonicotinoids, 869
organophosphates, 869
pyrethroid, 869
redbanded stink bug, 869
Dowling,
Aphis glycines, 648
electrical penetration graph, 648
host plant resistance, 648
soybean aphid, 648
downy brome,
BROTE, 348
Neoseiulus, 348
spider mites, 348
western corn rootworm, 348
drugstore beetle,
bed bug, 1636
cockroach, 1636
red flour beetle, 1636
rice weevil, 1636
dust,
economic cost, 2140
fumigant, 2140
soil barrier, 2140
termiticide, 2140

E

ear-feeding damage,
economic injury level, 2072
Euschistus servus, 2072
grain yield, 2072
kernel quality, 2072
eastern hemlock,
canopy translocation, 368
concentration, 368
hemlock woolly adelgid, 368
imidacloprid, 368
ecological negative cross-resistance,
Pectinophora gossypiella, 1821
resistance management, 1821
simulation modeling, 1821
Steinernema riobrave, 1821
economic cost,
dust, 2140
fumigant, 2140
soil barrier, 2140
termiticide, 2140
economic injury level,
ear-feeding damage, 2072
Euschistus servus, 2072
grain yield, 2072
kernel quality, 2072
ecotype,
Ostrinia nubilalis, 249
parasitoid, 249
synchronization, 249
Ectomyelois ceratoniae,
binomial sampling plan, 1474
Phoenix dactylifera, 1474
resampling validation, 1474
seasonal population dynamics, 1474
efficacy assessment,
infrared radiation, 1508
nonchemical method, 1508
Rhyzopertha dominica, 1508
egg laying,
acetamiprid, 379
Rhagoletis indifferens, 379
spinosad bait, 379
thiamethoxam, 379
electrical penetration graph,
Aphis glycines, 648
Dowling, 648
host plant resistance, 648
soybean aphid, 648
emamectin benzoate,
epigenetic transmission, 1347
Plutella xylostella, 1347

resistance, 1347
tolerance, 1347
emerald ash borer, 108, 597, 708, 1235, 1569
Asian longhorned beetle, 1569
azadirachtin, 708
biological invasions, 1569
firewood, 597, 1569
foliar uptake, 708
invasive species, 108
invasive species management, 1569
larval mortality, 708
nonindigenous species, 108
pest quarantines, 597
phytosanitation, 597
prism trap, 1235
regulatory compliance, 597
systemic injection, 708
trap height, 1235
urban forest, 108
visual cues, 1235
emergence,
low-temperature tolerance, 234
postemergence longevity, 234
stress, 234
emergence period,
emergence rate, 2028
humidity, 2028
Plutella xylostella, 2028
temperature, 2028
emergence rate,
emergence period, 2028
humidity, 2028
Plutella xylostella, 2028
temperature, 2028
emergence trap,
blueberries, 1258
Dasineura oxycoccana, 1258
distribution, 1258
rabbiteye, 1258
endoparasitoids,
biocontrol, 908
cytochrome oxidase subunit I gene, 908
DNA fingerprinting, 908
molecular markers, 908
endoplasmic reticulum,
methyl palmitate, 1372
mitochondria, 1372
Tetranychus viennensis, 1372
endosymbiont variation,
biotypes, 887
Diuraphis noxia, 887
entomopathogenic fungi,
biological control, 1094
defensive pheromone, 1094
solid phase microextraction, 1094
stored products pests, 1094
entomopathogenic nematodes, 1, 416, 1979
Acthina tumida, 1
Apis mellifera, 1
biological control, 416
Fraser fir, 1979
hazelnut, 416
Heterorhabditis spp., 1
nematode efficacy, 416
orchard floor cover, 416
Phyllophaga anxia, 1979
soil-applied insecticides, 1979
Steinernema spp., 1
Entostat powder,
autoconfusion, 2100
codlemone, 2100
flight tunnel, 2100
mating disruption, 2100
epidemiology,
Apis mellifera, 1517
classification and regression tree analysis, 1517
colony collapse disorder, 1517
pathogens, 1517
epigenetic transmission,
emamectin benzoate, 1347
Plutella xylostella, 1347
resistance, 1347
tolerance, 1347
epizootic,
antifungal, 1327
cellular encapsulation, 1327
grooming, 1327
Isoptera, 1327
eradication, 356, 1135
acoustic detection, 1135
Anthonomus grandis grandis, 356
boll weevil, 356
grandlure, 356
invasive species, 1135
trapping, 356
essential oil,
botanical acaricide, 1293
natural fumigant, 1293
Neoseiulus californicus, 1293
Tetranychus urticae, 1293
essential oils, 448, 781
Blattella germanica, 448, 781
fumigation, 781
ootheca, 448, 781
toxicity, 448
ethidium monoazide,
bacterial diversity, 1832
colonization, 1832
Musca domestica, 1832
16S rDNA, 1832
Ethovision,
image analysis, 219
IPM, 219
spray coverage, 219
water-sensitive spray cards, 219
Eucalyptus,
adventive range, 1943
insect herbivores, 1943
intentional introductions, 1943
invasive species, 1943
European corn borer, 1803, 2080
Bt maize, 1803
field corn, 2080
HDR strategy, 1803
infestation technique, 2080
Mon810, 1803
Ostrinia nubilalis, 2080
resistance management, 1803
Euschistus servus,
ear-feeding damage, 2072
economic injury level, 2072
grain yield, 2072
kernel quality, 2072
exotic, 1080, 1682
classical biological control, 1080
concealed host, 1080
firewood, 1682
forest pest, 1080
fuelwood, 1682
invasive, 1682
non-native, 1080
pathway, 1682
exotic species,
bark beetles, 1187
Ohio, 1187
phenology, 1187
semiochemicals, 1187
exotic species survey,
canopy malaise, 698
species richness, 698
trapping, 698
exposure period,
chlorantraniliprole, 1277
Choristoneura rosaceana, 1277
residues, 1277
spinetoram, 1277
external lesions,
boll damage, 2236
cotton, 2236
sampling, 2236
stink bug, 2236
extrafloral nectar,
apple, 1657
conservation biological control, 1657
Goniozus floridanus, 1657
Platynota idaeusalis, 1657

F

F_2 and F_1 screens,
Bt resistance, 472
Helicoverpa armigera, 472
fall armyworm,
Cry1F, 1031
transgenic maize, 1031
false codling moth,
CATTS, 265
controlled atmosphere, 265
heat, 265
postharvest, 265
fecundity, 292, 443, 662
alfalfa weevil, 292
development, 662
fertility, 662
life table entropy, 292
longevity, 443
methoxyfenozide, 662
modeling, 292
nymphal development, 443
Spodoptera exigua, 662
survivorship, 292
Symploce pallens, 443
feeding,
developmental period, 1744
Frankliniella occidentalis, 1744
oviposition, 1744
survival rate, 1744
feeding stimulant,
foraging, 2126
population, 2126
Reticulitermes flavipes, 2126
sugar, 2126

fertility,
development, 662
fecundity, 662
methoxyfenozide, 662
Spodoptera exigua, 662
field corn,
European corn borer, 2080
infestation technique, 2080
Ostrinia nubilalis, 2080
field experiments,
Aphis glycines, 1483
host plant resistance, 1483
predation, 1483
sampling, 1483
field-collected populations,
chlorantraniliprole, 140
insecticide resistance, 140
obliquebanded leafroller, 140
spinetoram, 140
fipronil, 433, 1761
California least tern, 1761
chemical barrier, 433
horizontal transfer, 1761
imidacloprid, 433
Linepithema humile, 1761
soil penetration, 433
termites, 433
firewood, 597, 1569, 1682
Asian longhorned beetle, 1569
biological invasions, 1569
emerald ash borer, 597, 1569
exotic, 1682
fuelwood, 1682
invasive, 1682
invasive species management, 1569
pathway, 1682
pest quarantines, 597
phytosanitation, 597
regulatory compliance, 597
fitness benefit,
Chrsoperla carnea, 823
organophosphate, 823
pyrethroids, 823
resistance, 823
fitness costs, 835, 1386
Bt resistance, 835
Cry1Ab, 1386
Cry2Ab, 835
Helicoverpa armigera, 835
high-dose refuge, 1386
resistance and insect resistance management, 1386
flavor,
Amegilla, 1286
horticulture, 1286
pollination, 1286
seeds, 1286
flight pattern,
aggregation, 95
Pinus yunnanensis, 95
Tomicus yunnanensis, 95
flight tunnel,
autoconfusion, 2100
codlemone, 2100
Entostat powder, 2100
mating disruption, 2100
flightless,
artificial selection, 1564
biological control, 1564
lady beetle, 1564
locomotor activity, 1564
Florida,
Anastrepha suspensa, 2214
host race formation, 2214
invasive insects, 2214
migration patterns, 2214
flour mill, 991, 1002
fumigation, 991, 1002
methyl bromide, 991
monitoring, 991
pheromone trapping, 1002
population dynamics, 1002
Tribolium castaneum, 991, 1002
Fluon,
Cerambycidae, 641
pheromone, 641
trap, 641
wood-boring insect, 641
foliar uptake,
azadirachtin, 708
emerald ash borer, 708
larval mortality, 708
systemic injection, 708
food consumption,
Coptotermes formosanus, 2132
naphthalene derivatives, 2132
repellency, 2132
foraging,
feeding stimulant, 2126
population, 2126
Reticulitermes flavipes, 2126
sugar, 2126
foraging territory,
Lincoln index, 2112
mark–release–recapture, 2112
Microcerotermes diversus, 2112
weighted mean model, 2112
forest pest,
classical biological control, 1080
concealed host, 1080
exotic, 1080
non-native, 1080
formulation,
Blattella germanica, 2118
microencapsulation, 2118
repellency, 2118
toxicity, 2118
Frankliniella occidentalis,
developmental period, 1744
feeding, 1744
oviposition, 1744
survival rate, 1744
Fraser fir,
entomopathogenic nematodes, 1979
Phyllophaga anxia, 1979
soil-applied insecticides, 1979
Fraxinus, 119, 603
Agrilus planipennis, 119
Agrilus, 603
borers, 119
fumigation, 603
imidacloprid, 119
quarantine, 603
sulfuryl fluoride, 603
freeze,
Aprostocetus hagenowii, 1770
heat, 1770
parasitoid, 1770
Periplaneta americana, 1770
French Guiana,
Anopheles spp., 973
collection methods, 973
fuelwood,
exotic, 1682
firewood, 1682
invasive, 1682
pathway, 1682
fumigant, 1988, 2140
Carposina niponensis, 1988
dust, 2140
economic cost, 2140
phosphine, 1988
soil barrier, 2140
termiticide, 2140
toxicity, 1988
fumigation, 603, 781, 991, 1002
Agrilus, 603
Blattella germanica, 781
essential oils, 781
flour mill, 991, 1002
Fraxinus, 603
methyl bromide, 991
monitoring, 991
ootheca, 781
pheromone trapping, 1002
population dynamics, 1002
quarantine, 603
sulfuryl fluoride, 603
Tribolium castaneum, 991, 1002
fungicide,
Listronotus maculicollis, 1810
P450 resistance, 1810
plant growth regulators, 1810
synergist, 1810

G

GABA receptor,
acaricide resistance, 1355
dieldrin, 1355
Rhipicephalus microplus, 1355
gamma radiation,
Cylas formicarius elegantulus, 2034
mating behavior, 2034
sterile insect technique (SIT), 2034
substerilization, 2034
gene flow,
genetics, 1012
isozyme, 1012
Stomoxys calcitrans, 1012
Thailand, 1012
genetic differentiation,
mitochondrial cytochrome oxidase subunit I, 2204
nucleotide diversity, 2204
Panonychus citri, 2204
population structure, 2204
genetic diversity,
biotype, 1454
microsatellite, 1454
Schizaphis graminum, 1454
simple sequence repeat, 1454
genetic markers,
Bactrocera jarvisi, 1071
Bactrocera tryoni, 1071
interspecific hybridization, 1071

genetic stock identification,
honey bee, 917
microsatellite, 917
single-nucleotide polymorphism, 917
genetics,
gene flow, 1012
isozyme, 1012
Stomoxys calcitrans, 1012
Thailand, 1012
geostatistics,
contour analysis, 1886
Mediterranean fruit fly, 1886
spatial statistics, 1886
variogram, 1886
global sampling,
mites, 2242
population dynamics, 2242
powdery mildew, 2242
thrips, 2242
glucosinolates,
artificial infestation, 1866
Brassica, 1866
Mamestra brassicae, 1866
plant defense, 1866
glutathione S-transferase,
B biotype, 158
Bemisia tabaci, 158
cytochrome P450, 158
monooxygenase, 158
resistance to pyriproxyfen, 158
Glycine max,
antibiosis, 1405
Aphis glycines, 1405
host plant resistance, 1405
tolerance, 1405
Gnathotrichus materiarius,
conophthorin, 2094
Ips grandicollis, 2094
Ips pini, 2094
Xylosandrus germanus, 2094
Goniozus floridanus,
apple, 1657
conservation biological control, 1657
extrafloral nectar, 1657
Platynota idaeusalis, 1657
grain yield,
ear-feeding damage, 2072
economic injury level, 2072
Euschistus servus, 2072
kernel quality, 2072
grandisoic acid,
benzaldehyde, 744
Circle trap, 744
plum curculio, 744
pyramid trap, 744
grandlure,
Anthonomus grandis grandis, 356
boll weevil, 356
eradication, 356
trapping, 356
granulovirus proteins,
Helicoverpa armigera, 257
Mamestra brassicae, 257, 257
nucleopolyhedrovirus, 257
viral enhancement, 257
grape,
alternate host plants, 1051
citrus, 1051
Pierce's disease, 1051
vector control, 1051
grape benchgraft,
controlled atmosphere, 272
mealybug, 272
Planococcus ficus, 272
ultralow oxygen, 272
grapefruit-lemon,
canopy distribution, 47
colored sticky cards, 47
host plants, 47
Kelly's citrus thrips, 47
grapevine,
citrus mealybug, 619
Grapevine leafroll-associated virus 3, 619
insect, 619
Planococcus citri, 619
Grapevine leafroll-associated virus 3,
citrus mealybug, 619
grapevine, 619
insect, 619
Planococcus citri, 619
Grapholita molesta, 1299, 1815
acetamiprid, 1815
chlorantraniliprole, 1815
mating disruption, 1299
paraffin wax., 1299
pheromone dispenser, 1299
point source, 1299
spinetoram, 1815
spinosad, 1815
greenhouse tomato,
Aculops lycopersici, 563
Amblyseius swirskii, 563
biological control, 563
grooming,
antifungal, 1327
cellular encapsulation, 1327
epizootic, 1327
Isoptera, 1327
growth and development,
insect growth regulator, 340
lethal and sublethal effects, 340
obliquebanded leafroller, 340
pyriproxyfen, 340
growth and reproduction,
artificial diet, 17
OB yield, 17
SeMNPV, 17
Spodoptera exigua, 17

H

haplotypes,
migration, 360
Spodoptera frugiperda, 360
Hass avocado,
Anastrepha ludens, 2000
cold treatment, 2000
risk analysis, 2000
Tephritidae, 2000
hazelnut,
biological control, 416
entomopathogenic nematodes, 416
nematode efficacy, 416
orchard floor cover, 416
HDR strategy,
Bt maize, 1803
European corn borer, 1803
Mon810, 1803
resistance management, 1803
heat, 265, 1770, 1909
Anobiidae, 1909
Aprostocetus hagenowii, 1770
CATTS, 265
cold, 1909
controlled atmosphere, 265
false codling moth, 265
freeze, 1770
parasitoid, 1770
Periplaneta americana, 1770
postharvest, 265
supercooling point, 1909
tolerance, 1909
heat unit,
Acrobasis nuxvorella, 735
biofix, 735
Carya, 735
physiological model, 735
Helicoverpa armigera, 257, 472, 835, 1420, 2169
Bt resistance, 472, 835
cross-tolerance, 2169
Cry1Ac, 2169
Cry2Ab, 835
demographic parameters, 1420
F_2 and F_1 screens, 472
fitness costs, 835
granulovirus proteins, 257
Mamestra brassicae, 257, 257
nucleopolyhedrovirus, 257
nutritional indices, 1420
soybean cultivars, 1420
Vip3Aa, 2169
viral enhancement, 257
Helicoverpa punctigera,
Bacillus thuringiensis, 2147
Cry1Ac, 2147
Cry2Ab, 2147
resistance, 2147
Hemiberlesia lanatiae,
Abgrallaspis aguacatae, 1172
Aspidiotus nerii, 1172
Diaspidiotus perniciosus, 1172
phoresy, 1172
hemlock woolly adelgid,
canopy translocation, 368
concentration, 368
eastern hemlock, 368
imidacloprid, 368
herbivory,
biotic, 516
phosphorus deficiency, 516
stem mining, 516
water stress, 516
heritability,
chlorantraniliprole, 1378
obliquebanded leafroller, 1378
resistance, 1378
spinetoram, 1378
Hessian fly genotypes,
Mayetiola destructor, 2229
plant resistance, 2229
wheat breeding, 2229
Heterorhabditis bacteriophora,
Rhyzopertha dominica, 977
Steinernema carpocapsae, 977
Steinernema feltiae, 977
stored product pests *kuehniella*, 977

Heterorhabditis spp.,
Aethina tumida, 1
Apis mellifera, 1
entomopathogenic nematodes, 1
Steinernema spp., 1
high-dose refuge,
Cry1Ab, 1386
fitness costs, 1386
resistance and insect resistance management, 1386
honey bee, 10, 917, 1039
Aethina tumida, 10
Apis mellifera, 10
genetic stock identification, 917
microsatellite, 917
Noseme ceranae, 10
parasitic mite, 1039
pest management, 1039
single-nucleotide polymorphism, 917
Varroa destructor, 10
horizontal toxicity,
barrier, 754
insecticide, 754
Solenopsis invicta, 754
horizontal transfer,
California least tern, 1761
fipronil, 1761
Linepithema humile, 1761
horn fly,
Bacillus thuringiensis, 1019
microbial control, 1019
stable fly, 1019
horticulture,
Amegilla, 1286
flavor, 1286
pollination, 1286
seeds, 1286
host plant resistance, 648, 1405, 1483, 1860
adult damage, 1860
antibiosis, 1405
Aphis glycines, 648, 1405, 1483
Brachiaria, 1860
Dowling, 648
electrical penetration graph, 648
field experiments, 1483
Glycine max, 1405
life stages, 1860
predation, 1483
sampling, 1483
soybean aphid, 648
spittlebugs, 1860
tolerance, 1405
host plant selection,
mechanism, 1180
oviposition, 1180
preference, 1180
resistance, 1180
host plants,
canopy distribution, 47
colored sticky cards, 47
grapefruit-lemon, 47
Kelly's citrus thrips, 47
host preference,
Africa, 70
Bactrocera latifrons, 70
host range, 70
Solanaceae, 70
host race formation,
Anastrepha suspensa, 2214
Florida, 2214
invasive insects, 2214
migration patterns, 2214
host range,
Africa, 70
Bactrocera latifrons, 70
host preference, 70
Solanaceae, 70
host suitability,
alternate host, 2009
life table, 2009
Lygus hesperus, 2009
western tarnished plant bug, 2009
house fly, 550, 1929
imidacloprid, 550
metaflumizone, 550
Musca domestica, 550
muscalure, 1929
spinosad, 550
(Z)-9-tricosene, 1929
huanglongbing, 127, 541, 1531, 1972
abandoned citrus, 1972
acquisition, 1531
Asian citrus psyllid, 1972
Candidatus Liberibacter asiaticus, 1972
citrus greening, 127, 1972
citrus greening disease, 541, 1531
inoculation, 1531
king mandarin, 127
monitoring, 541
population, 127
systemic insecticide, 127
vector, 1531
humidity,
emergence period, 2028
emergence rate, 2028
Plutella xylostella, 2028
temperature, 2028
hydrolyzed yeast,
Caribbean fruit fly, 40
male competitiveness, 40
methoprene, 40
sexual performance, 40
Hypothenemus hampei,
Coffea arabica, 1159
infestation, 1159
reproduction, 1159
temperature, 1159

I

image analysis, 219, 1464
crop rotation, 1464
Ethovision, 219
IPM, 219
soybean herbivory, 1464
spray coverage, 219
water-sensitive spray cards, 219
western corn rootworm, 1464
imidacloprid, 119, 368, 433, 550, 1711
Agrilus planipennis, 119
borers, 119
canopy translocation, 368
chemical barrier, 433
concentration, 368
control level, 1711
Diaphorina citri, 1711
eastern hemlock, 368
fipronil, 433
Fraxinus, 119
hemlock woolly adelgid, 368
house fly, 550
metaflumizone, 550
Musca domestica, 550
Phyllocnistis citrella, 1711
soil penetration, 433
spinosad, 550
systemic uptake, 1711
termites, 433
Indianmeal moth,
product moisture, 612
relative humidity, 612
vacuum treatment, 612
indoxacarb,
Reticulitermes flavipes, 762
sublethal behaviors, 762
Inesfly IGR FITO,
palm, 402
Phoenix canariensis, 402
Rhynchophorus ferrugineus, 402
infection frequency,
Bemisia tabaci, 1848
parasitoid, 1848
predator, 1848
Wolbachia supergroups, 1848
infestation,
Coffea arabica, 1159
Hypothenemus hampei, 1159
reproduction, 1159
temperature, 1159
infestation technique,
European corn borer, 2080
field corn, 2080
Ostrinia nubilalis, 2080
inflorescence,
Mangifera indica, 631
nondestructive sampling methods, 631
sticky trap, 631
Thripidae, 631
infrared radiation,
efficacy assessment, 1508
nonchemical method, 1508
Rhyzopertha dominica, 1508
inoculation,
acquisition, 1531
citrus greening disease, 1531
huanglongbing, 1531
vector, 1531
insect,
citrus mealybug, 619
grapevine, 619
Grapevine leafroll-associated virus 3, 619
Planococcus citri, 619
insect behavior,
Diabrotica virgifera virgifera, 2052
larvae, 2052
resistance, 2052
Zea mays, 2052
insect diet,
Blissus insularis, 726
chinch bug, 726
rearing methods, 726
insect growth regulator, 340, 2174
Bemisia tabaci, 2174
growth and development, 340
insecticide resistance, 2174

lethal and sublethal effects, 340
neonicotinoids, 2174
obliquebanded leafroller, 340
pyriproxyfen, 340
Q biotype, 2174
insect herbivores,
adventive range, 1943
Eucalyptus, 1943
intentional introductions, 1943
invasive species, 1943
insect management,
IPM, 1896
stochastic dynamic programming, 1896
stored corn, 1896
insect population,
mating disruption, 1621
olive yield, 1621
Zeuzera pyrina, 1621
insect resistance evolution,
Bt maize, 1394
Cry1Ab, 1394
Cry3Bb, 1394
transgenic maize, 1394
insecticide, 308, 668, 754, 1315
action threshold, 1315
Allium cepa, 1315
Apoidea, 668
barrier, 754
biological control, 308
horizontal toxicity, 754
nontarget effects, 668
Solenopsis invicta, 754
sublethal effects, 308
Thrips tabaci, 1315
Trichogramma evanescens, 308
Trichogramma semblidis, 308
Vaccinium corymbosum, 668
wild bee conservation, 668
insecticide application timing,
desiccation, 1306
integrated pest management, 1306
irrigation, 1306
Phthorimaea operculella, 1306
insecticide bait,
dairy pest management, 1937
insecticide resistance, 1937
integrated pest management, 1937
Musca domestica, 1937
insecticide residue,
acetamiprid, 2107
Contarinia nasturtii, 2107
invasive species, 2107
spray coverage, 2107
insecticide resistance, 140, 374, 460, 816, 1365, 1400, 1937, 2174
abamectin susceptibility, 816
aphids, 374
Aphis pomi, 374
Aphis spiraecola, 374
Bemisia tabaci, 2174
bioassays, 816
Blattella germanica, 460
chlorantraniliprole, 140
citrus, 1400
dairy pest management, 1937
field-collected populations, 140
insect growth regulators, 2174
insecticide bait, 1937
integrated pest management, 1937
LD-P line, 1365
Leptinotarsa decemlineata, 1365
Musca domestica, 1937
neonicotinoids, 2174
obliquebanded leafroller, 140
organophosphate, 1400
pear psylla, 816
pest control, 816
Q biotype, 2174
resistance mechanism, 460
Singapore, 460
spinetoram, 140
susceptibility, 374
synergism, 460
synergist, 1365
topical bioassay, 1365
insecticide resistance management,
reproductive biology, 861
sperm precedence, 861
tobacco budworm, 861
YHD2, 861
insecticide resistance susceptibility,
adults, 2164
larvae, 2164
Thysanoptera, 2164
integrated pest management, 386, 1306, 1937
coverage, 386
dairy pest management, 1937
desiccation, 1306
insecticide application timing, 1306
insecticide bait, 1937
insecticide resistance, 1937
irrigation, 1306
mineral oil, 386
Musca domestica, 1937
Phthorimaea operculella, 1306
twospotted spider mite, 386
integrated pest management in viticulture,
biological control, 2061
Calepitrimerus vitis, 2061
Colomerus vitis, 2061
Phytoseiidae, 2061
intentional introductions,
adventive range, 1943
Eucalyptus, 1943
insect herbivores, 1943
invasive species, 1943
interspecific, 1754, 1775
aggression, 1775
agonism, 1754
Anoplolepis gracilipes, 1775
intraspecific, 1754, 1775
invasive species, 1775
Microcerotermes crassus, 1754
survival, 1754
interspecific hybridization,
Bactrocera jarvisi, 1071
Bactrocera tryoni, 1071
genetic markers, 1071
intraguild competition,
Cry1Ab, 54
MON810, 54
Striacosta albicosta, 54
YieldGard, 54
intraguild predation,
pest management tools, 1086
pollen, 1086
predators, 1086
western flower thrips, 1086
intraspecific, 1754, 1775
aggression, 1775
agonism, 1754
Anoplolepis gracilipes, 1775
interspecific, 1754, 1775
invasive species, 1775
Microcerotermes crassus, 1754
survival, 1754
intrinsic rate of increase,
Melanaphis sacchari, 1431
Saccharum resistance, 1431
Sipha flava, 1431
inundative release,
Aphthona, 242
biological control, 242
leafy spurge, 242
riparian area, 242
invasive,
exotic, 1682
firewood, 1682
fuelwood, 1682
pathway, 1682
invasive fruit flies,
male lures, 2042
mark–recapture, 2042
surveillance, 2042
invasive insects,
Anastrepha suspensa, 2214
Florida, 2214
host race formation, 2214
migration patterns, 2214
invasive species, 108, 1135, 1775, 1875, 1943, 2107
acetamiprid, 2107
acoustic detection, 1135
adventive range, 1943
aggression, 1775
Anoplolepis gracilipes, 1775
Contarinia nasturtii, 2107
Diuraphis frequens, 1875
Diuraphis mexicana, 1875
Diuraphis noxia, 1875
Diuraphis tritici, 1875
emerald ash borer, 108
eradication, 1135
Eucalyptus, 1943
insect herbivores, 1943
insecticide residue, 2107
intentional introductions, 1943
interspecific, 1775
intraspecific, 1775
nonindigenous species, 108
spray coverage, 2107
urban forest, 108
invasive species management,
Asian longhorned beetle, 1569
biological invasions, 1569
emerald ash borer, 1569
firewood, 1569
IPM, 219, 654, 1735, 1896
armored scales, 1735
bark absorption, 1735
canola, 654
Christmas tree, 1735
Ethovision, 219
image analysis, 219
insect management, 1896
Lepidoptera, 654

pheromone monitoring, 654
Plutella xylostella, 654
spray coverage, 219
stochastic dynamic programming, 1896
stored corn, 1896
systemic insecticide, 1735
water-sensitive spray cards, 219
Ipomoea batatas,
Bt bioassays, 1493
Bt toxins, 1493
Ips grandicollis,
conophthorin, 2094
Gnathotrichus materiarius, 2094
Ips pini, 2094
Xylosandrus germanus, 2094
Ips pini, 1693, 2094
conophthorin, 2094
Dendroctonus, 1693
Gnathotrichus materiarius, 2094
Ips grandicollis, 2094
microclimate, 1693
Pinus ponderosa, 1693
thinning, 1693
Xylosandrus germanus, 2094
Iris yellow spot virus,
Allium cepa, 925
onion resistance, 925
Thrips tabaci, 925
irradiation, 85, 1950
Alpha radiation, 85
Anastrepha fraterculus, 85
Ceratitis capitata, 85
commodity treatment, 1950
disinfestation, 1950
quarantine, 1950
X radiation, 85
irrigation,
desiccation, 1306
insecticide application timing, 1306
integrated pest management, 1306
Phthorimaea operculella, 1306
isolation-by-distance,
amplified fragment length polymorphism, 958
biogeography, 958
biotypes, 958
Diuraphis noxia, 958
Isoptera,
antifungal, 1327
cellular encapsulation, 1327
epizootic, 1327
grooming, 1327
isozyme,
gene flow, 1012
genetics, 1012
Stomoxys calcitrans, 1012
Thailand, 1012
ITS,
mealybug, 25
molecular identification, 25
pear, 25
species-specific primer, 25

J

juvenile hormone,
cannibalism, 577
nucleopolyhedrovirus production, 577
Spodoptera exigua, 577

K

kaolin,
apple maggot, 394
behavior, 394
particle film technology, 394
Surround WP, 394
Kelly's citrus thrips,
canopy distribution, 47
colored sticky cards, 47
grapefruit-lemon, 47
host plants, 47
Kelly's citrus thrips, 47
kernel quality,
ear-feeding damage, 2072
economic injury level, 2072
Euschistus servus, 2072
grain yield, 2072
king mandarin,
citrus greening, 127
Huanglongbing, 127
population, 127
systemic insecticide, 127

L

laboratory adaptation,
Coleoptera, 1144
Cylas formicarius elegantulus, 1144
sterile insect technique, 1144
lady beetle,
artificial selection, 1564
biological control, 1564
flightless, 1564
locomotor activity, 1564
landscape,
Leptinotarsa decemlineata, 1613
relative humidity, 1613
solar radiation, 1613
temperature, 1613
Laricobius nigrinus,
nontarget, 1222
prey-mediated effects, 1222
Sasajiscymnus tsugae, 1222
Tsuga canadensis, 1222
larvae, 2052, 2164
adults, 2164
Diabrotica virgifera virgifera, 2052
insect behavior, 2052
insecticide resistance susceptibility, 2164
resistance, 2052
Thysanoptera, 2164
Zea mays, 2052
larval mortality,
azadirachtin, 708
emerald ash borer, 708
foliar uptake, 708
systemic injection, 708
laundry,
bed bug, 136
Cimex lectularius, 136
control, 136
LD-P line,
insecticide resistance, 1365
Leptinotarsa decemlineata, 1365
synergist, 1365
topical bioassay, 1365
leafy spurge,
Aphthona, 242
biological control, 242
inundative release, 242
riparian area, 242
Lepidoptera,
canola, 654
IPM, 654
pheromone monitoring, 654
Plutella xylostella, 654
Lepinotus reticulatus,
Rhyzopertha dominica, 197
Sitophilus oryzae, 197
stored products, 197
Tribolium castaneum, 197
Leptinotarsa decemlineata, 1365, 1613
insecticide resistance, 1365
landscape, 1613
LD-P line, 1365
relative humidity, 1613
solar radiation, 1613
synergist, 1365
temperature, 1613
topical bioassay, 1365
lethal and sublethal effects,
growth and development, 340
insect growth regulator, 340
obliquebanded leafroller, 340
pyriproxyfen, 340
liberibacter,
carrot, 1060
plant pathogen, 1060
psyllid, 1060
Trioza apicalis, 1060
life stages,
adult damage, 1860
Brachiaria, 1860
host plant resistance, 1860
spittlebugs, 1860
life table, 1628, 1729, 2009, 2223
alternate host, 2009
Bactericera cockerelli, 1729
Chaitophorus leucomelas, 2223
host suitability, 2009
Lygus hesperus, 2009
mortality, 1729
Nilaparvata lugens, 1628
O. officinalis, 1628
O. rufipogon, 1628
population dynamics, 1729
Populus, 2223
susceptibility, 2223
western tarnished plant bug, 2009
Zebra Chip, 1729
life table entropy,
alfalfa weevil, 292
fecundity, 292
modeling, 292
survivorship, 292
life-stage activity,
Acrobasis vaccinii, 1720
parasitoid, 1720
reduced-risk insecticide, 1720
life-table analysis,
canola cultivars, 2019
model based survival, 2019
Plutella xylostella, 2019
Lincoln index,
foraging territory, 2112
mark–release–recapture, 2112
Microcerotermes diversus, 2112
weighted mean model, 2112

Linepithema humile, 1761, 1784, 1790
Argentine ant, 1784
California least tern, 1761
Coccidae, 1790
control, 1784
fipronil, 1761
horizontal transfer, 1761
Mesolecanium nigrofasciatum, 1790
pest management, 1790
red maple, 1790
synthetic trail pheromone, 1784
(Z)-9-hexadecenal, 1784
Liposcelis,
varietal resistance, 985
wheat classes, 985
Liriomyza trifolii,
abamectin, 2197
cyromazine, 2197
random amplified polymorphic DNA polymerase chain reaction, 2197
resistance, 2197
Listronotus maculicollis,
fungicide, 1810
P450 resistance, 1810
plant growth regulators, 1810
synergist, 1810
locomotor activity,
artificial selection, 1564
biological control, 1564
flightless, 1564
lady beetle, 1564
longevity,
fecundity, 443
nymphal development, 443
Symploce pallens, 443
low-temperature tolerance,
emergence, 234
postemergence longevity, 234
stress, 234
Loxosceles, 166, 808
Arachnida, 808
brown recluse spider, 808
loxoscelism, 166
pest control, 166
pyrethroids, 166
sericophily, 808
Sicariidae, 166
urban entomology, 808
loxoscelism,
Loxosceles, 166
pest control, 166
pyrethroids, 166
Sicariidae, 166
Lygus hesperus,
alternate host, 2009
host suitability, 2009
life table, 2009
western tarnished plant bug, 2009
Lymantria dispar,
biomass production, 718
microbial pesticides, 718
Populus, 718
short rotation trees, 718

M

male annihilation treatment,
Bactrocera cucurbitae, 1594
control, 1594
Min-U-Gel, 1594
naled, 1594
male competitiveness,
Caribbean fruit fly, 40
hydrolyzed yeast, 40
methoprene, 40
sexual performance, 40
male lures,
invasive fruit flies, 2042
mark–recapture, 2042
surveillance, 2042
Malpighian tubules,
ant bait, 676
microscopy, 676
midgut, 676
postpharyngeal gland, 676
Mamestra brassicae, 257, 257, 1866
artificial infestation, 1866
Brassica, 1866
glucosinolates, 1866
granulovirus proteins, 257, 257
Helicoverpa armigera, 257, 257
nucleopolyhedrovirus, 257, 257
plant defense, 1866
viral enhancement, 257, 257
Mangifera indica,
inflorescence, 631
nondestructive sampling methods, 631
sticky trap, 631
Thripidae, 631
mark–recapture,
invasive fruit flies, 2042
male lures, 2042
mark–recapture, 2042
surveillance, 2042
mark–release–recapture, 1242, 2112
attractive range, 1242
codling moth, 1242
foraging territory, 2112
Lincoln index, 2112
mark–release–recapture, 1242, 2112
mating disruption, 1242
Microcerotermes diversus, 2112
spatial scale, 1242
weighted mean model, 2112
mating behavior,
Cylas formicarius elegantulus, 2034
gamma radiation, 2034
sterile insect technique (SIT), 2034
substerilization, 2034
mating disruption, 1242, 1299, 1621, 2100
attractive range, 1242
autoconfusion, 2100
codlemone, 2100
codling moth, 1242
Entostat powder, 2100
flight tunnel, 2100
Grapholita molesta, 1299
insect population, 1621
mark–release–recapture, 1242
olive yield, 1621
paraffin wax., 1299
pheromone dispenser, 1299
point source, 1299
spatial scale, 1242
Zeuzera pyrina, 1621
Mayetiola destructor, 178, 2229
Hessian fly genotypes, 2229
phytohormones, 178
plant resistance, 2229
resistance, 178
wheat, 178
wheat breeding, 2229
mCry3A,
Bacillus thuringiensis, 2187
Bt, 2187
Diabrotica virgifera virgifera, 2187
resistance management, 2187
mealybug, 25, 272
controlled atmosphere, 272
grape benchgraft, 272
ITS, 25
molecular identification, 25
pear, 25
Planococcus ficus, 272
species-specific primer, 25
ultralow oxygen, 272
mechanism,
host plant selection, 1180
oviposition, 1180
preference, 1180
resistance, 1180
Mediterranean fruit fly, 1586, 1886
attractants, 1586
ceralure, 1586
contour analysis, 1886
detection, 1586
geostatistics, 1886
spatial statistics, 1886
trimedlure, 1586
variogram, 1886
Melanaphis sacchari,
intrinsic rate of increase, 1431
Saccharum resistance, 1431
Sipha flava, 1431
Melanotus communis,
Conoderus spp., 2087
crop rotation, 2087
wireworm, 2087
Mesolecanium nigrofasciatum,
Coccidae, 1790
Linepithema humile, 1790
pest management, 1790
red maple, 1790
metaflumizone,
house fly, 550
imidacloprid, 550
Musca domestica, 550
spinosad, 550
methoprene,
Caribbean fruit fly, 40
hydrolyzed yeast, 40
male competitiveness, 40
sexual performance, 40
methoxyfenozide, 662, 1704
codlemone, 1704
Cydia pomonella, 1704
development, 662
fecundity, 662
fertility, 662
responsiveness, 1704
Spodoptera exigua, 662
sublethal effects, 1704
methyl (2*E*6*E*)-farnesoate,
aggregation pheromone, 1603
methyl (2*E*6*E*)-farnesoate, 1603

methyl (*E*)-5-2610-trimethyl-59-undecadienoate, 1603
methyl (*E*)-6-23-dihydrofarnesoate, 1603
methyl geranate, 1603
methyl (*E*)-5-2610-trimethyl-59-undecadienoate,
aggregation pheromone, 1603
methyl (2*E*6*E*)-farnesoate, 1603
methyl (*E*)-5-2610-trimethyl-59-undecadienoate, 1603
methyl (*E*)-6-23-dihydrofarnesoate, 1603
methyl geranate, 1603
methyl (*E*)-6-23-dihydrofarnesoate,
aggregation pheromone, 1603
methyl (2*E*6*E*)-farnesoate, 1603
methyl (*E*)-5-2610-trimethyl-59-undecadienoate, 1603
methyl (*E*)-6-23-dihydrofarnesoate, 1603
methyl geranate, 1603
methyl bromide,
flour mill, 991
fumigation, 991
monitoring, 991
Tribolium castaneum, 991
methyl geranate,
aggregation pheromone, 1603
methyl (2*E*6*E*)-farnesoate, 1603
methyl (*E*)-5-2610-trimethyl-59-undecadienoate, 1603
methyl (*E*)-6-23-dihydrofarnesoate, 1603
methyl palmitate,
endoplasmic reticulum, 1372
mitochondria, 1372
Tetranychus viennensis, 1372
México,
aluminum-carboxymethylcellulose, 570
Bti, 570
microbial control, 570
microencapsulation, 570
microbial control, 570, 1019, 1555
Agrotis ipsilon, 1555
Agrotis ipsilon nucleopolyhedrovirus, 1555
aluminum-carboxymethylcellulose, 570
Bacillus thuringiensis, 1019
baculovirus, 1555
Bti, 570
horn fly, 1019
México, 570
microencapsulation, 570
natural enemies, 1555
stable fly, 1019
microbial pesticides,
biomass production, 718
Lymantria dispar, 718
Populus, 718
short rotation trees, 718
Microcerotermes crassus, 437, 1754
agonism, 1754
Coptotermes gestroi, 437
interspecific, 1754
intraspecific, 1754
moisture, 437
survival, 1754
tunneling activity, 437
wood consumption, 437
Microcerotermes diversus,
foraging territory, 2112
Lincoln index, 2112
mark–release–recapture, 2112
weighted mean model, 2112
microclimate,
Dendroctonus, 1693
Ips pini, 1693
Pinus ponderosa, 1693
thinning, 1693
microencapsulation, 570, 2118
aluminum-carboxymethylcellulose, 570
Blattella germanica, 2118
Bti, 570
formulation, 2118
México, 570
microbial control, 570
repellency, 2118
toxicity, 2118
microsatellite, 791, 917, 1454
biotype, 1454
colony elimination, 791
colony suppression, 791
genetic diversity, 1454
genetic stock identification, 917
honey bee, 917
Reticulitermes flavipes, 791
Reticulitermes virginicus, 791
Schizaphis graminum, 1454
simple sequence repeat, 1454
single-nucleotide polymorphism, 917
microscopy,
ant bait, 676
Malpighian tubules, 676
midgut, 676
postpharyngeal gland, 676
midgut,
ant bait, 676
Malpighian tubules, 676
microscopy, 676
postpharyngeal gland, 676
migration, 186, 360
Aphis glycines, 186
haplotypes, 360
sampling procedure, 186
soybean aphid, 186
Spodoptera frugiperda, 360
suction trap, 186
migration patterns,
Anastrepha suspensa, 2214
Florida, 2214
host race formation, 2214
invasive insects, 2214
Min-U-Gel,
Bactrocera cucurbitae, 1594
control, 1594
male annihilation treatment, 1594
naled, 1594
mineral oil,
coverage, 386
integrated pest management, 386
twospotted spider mite, 386
mites,
global sampling, 2242
population dynamics, 2242
powdery mildew, 2242
thrips, 2242
mitochondria,
endoplasmic reticulum, 1372
methyl palmitate, 1372
Tetranychus viennensis, 1372
mitochondrial cytochrome oxidase subunit I,
genetic differentiation, 2204
nucleotide diversity, 2204
Panonychus citri, 2204
population structure, 2204
mitochondrial DNA,
Ceratitis species, 1842
Dacus species, 1842
molecular taxonomy, 1842
model based survival,
canola cultivars, 2019
life-table analysis, 2019
Plutella xylostella, 2019
modeling,
alfalfa weevil, 292
fecundity, 292
life table entropy, 292
survivorship, 292
moisture,
Coptotermes gestroi, 437
Microcerotermes crassus, 437
tunneling activity, 437
wood consumption, 437
molecular diagnostic marker,
cytochrome oxidase subunit II, 898
pest management, 898
weevil, 898
molecular identification, 25, 877
barcoding, 877
cytochrome c oxidase I gene, 877
ITS, 25
mealybug, 25
pear, 25
28SD2, 877
species-specific primer, 25
tospovirus vector, 877
molecular markers,
biocontrol, 908
cytochrome oxidase subunit I gene, 908
DNA fingerprinting, 908
endoparasitoids, 908
molecular taxonomy,
Ceratitis species, 1842
Dacus species, 1842
mitochondrial DNA, 1842
MON810,
Cry1Ab, 54
intraguild competition, 54
Striacosta albicosta, 54
YieldGard, 54
Mon810,
Bt maize, 1803
European corn borer, 1803
HDR strategy, 1803
resistance management, 1803
monitoring, 172, 541, 991
bed bug, 172
Cimex lectularius, 172
citrus greening disease, 541
dispersal, 172
flour mill, 991
fumigation, 991

huanglongbing, 541
methyl bromide, 991
Tribolium castaneum, 991
monitoring insecticide resistance,
Apis mellifera, 1797
Varroa destructor, 1797
monooxygenase,
B biotype, 158
Bemisia tabaci, 158
cytochrome P450, 158
glutathione S-transferase, 158
resistance to pyriproxyfen, 158
mortality,
Bactericera cockerelli, 1729
life table, 1729
population dynamics, 1729
Zebra Chip, 1729
movement,
dispersal, 63
path, 63
tracking, 63
Musca domestica, 550, 1832, 1937
bacterial diversity, 1832
colonization, 1832
dairy pest management, 1937
ethidium monoazide, 1832
house fly, 550
imidacloprid, 550
insecticide bait, 1937
insecticide resistance, 1937
integrated pest management, 1937
metaflumizone, 550
16S rDNA, 1832
spinosad, 550
muscalure,
house fly, 1929
(Z)-9-tricosene, 1929

N

naled, 409, 1594
Bactrocera cucurbitae, 409, 1594
Bactrocera dorsalis, 409
control, 1594
male annihilation treatment, 1594
Min-U-Gel, 1594
spinosad, 409
tephritid flies, 409
naphthalene derivatives,
Coptotermes formosanus, 2132
food consumption, 2132
repellency, 2132
natural enemies,
Agrotis ipsilon, 1555
Agrotis ipsilon nucleopolyhedrovirus, 1555
baculovirus, 1555
microbial control, 1555
natural fumigant,
botanical acaricide, 1293
essential oil, 1293
Neoseiulus californicus, 1293
Tetranychus urticae, 1293
natural insecticide,
Brevicoryne brassicae, 492
camptothecin, 492
Chilo suppressalis, 492
Nilaparvata lugens, 492
navel orangeworm,
almonds, 1250
degree-days, 1250
development, 1250
survival, 1250
nematode efficacy,
biological control, 416
entomopathogenic nematodes, 416
hazelnut, 416
orchard floor cover, 416
neonicotinoids, 869, 2174
Bemisia tabaci, 2174
dose–mortality analysis, 869
insect growth regulators, 2174
insecticide resistance, 2174
organophosphates, 869
pyrethroid, 869
Q biotype, 2174
redbanded stink bug, 869
Neoseiulus californicus,
botanical acaricide, 1293
essential oil, 1293
natural fumigant, 1293
Tetranychus urticae, 1293
Neoseiulus,
BROTE, 348
downy brome, 348
spider mites, 348
western corn rootworm, 348
nest temperature,
baiting, 423
chitin synthesis inhibitor, 423
Coptotermes, 423
peritrophic membrane, 423
new host association,
Curculio caryae, 1312
quarantine threat, 1312
weevil DNA, 1312
Nilaparvata lugens, 492, 1628
Brevicoryne brassicae, 492
camptothecin, 492
Chilo suppressalis, 492
life table, 1628
natural insecticide, 492
O. officinalis, 1628
O. rufipogon, 1628
non-native,
classical biological control, 1080
concealed host, 1080
exotic, 1080
forest pest, 1080
nonchemical method,
efficacy assessment, 1508
infrared radiation, 1508
Rhyzopertha dominica, 1508
nondestructive sampling methods,
inflorescence, 631
Mangifera indica, 631
sticky trap, 631
Thripidae, 631
nonindigenous species,
emerald ash borer, 108
invasive species, 108
urban forest, 108
nonpersistent virus,
Aphis glycines, 1670
Phaseolus vulgaris, 1670
seasonal phenology, 1670
nontarget,
Laricobius nigrinus, 1222
prey-mediated effects, 1222
Sasajiscymnus tsugae, 1222
Tsuga canadensis, 1222
nontarget effects,
Apoidea, 668
insecticides, 668
Vaccinium corymbosum, 668
wild bee conservation, 668
Nosema ceranae,
Aethina tumida, 10
Apis mellifera, 10
honey bee, 10
Varroa destructor, 10
nucleopolyhedrovirus,
granulovirus proteins, 257
Helicoverpa armigera, 257
Mamestra brassicae, 257, 257
viral enhancement, 257
nucleopolyhedrovirus production,
cannibalism, 577
juvenile hormone, 577
Spodoptera exigua, 577
nucleotide diversity,
genetic differentiation, 2204
mitochondrial cytochrome oxidase subunit I, 2204
Panonychus citri, 2204
population structure, 2204
nutritional indices,
demographic parameters, 1420
Helicoverpa armigera, 1420
soybean cultivars, 1420
nymphal development,
fecundity, 443
longevity, 443
Symploce pallens, 443

O

O. officinalis,
life table, 1628
Nilaparvata lugens, 1628
O. rufipogon, 1628
O. rufipogon,
life table, 1628
Nilaparvata lugens, 1628
O. officinalis, 1628
OB yield,
artificial diet, 17
growth and reproduction, 17
SeMNPV, 17
Spodoptera exigua, 17
obliquebanded leafroller, 140, 340, 1378
chlorantraniliprole, 140, 1378
field-collected populations, 140
growth and development, 340
heritability, 1378
insect growth regulator, 340
insecticide resistance, 140
lethal and sublethal effects, 340
pyriproxyfen, 340
resistance, 1378
spinetoram, 140, 1378
Ohio,
bark beetles, 1187
exotic species, 1187
phenology, 1187
semiochemicals, 1187
olive yield,
insect population, 1621
mating disruption, 1621
Zeuzera pyrina, 1621

onion resistance,
Allium cepa, 925
Iris yellow spot virus, 925
Thrips tabaci, 925
ootheca, 448, 781
Blattella germanica, 448, 781
essential oils, 448, 781
fumigation, 781
toxicity, 448
orchard floor cover,
biological control, 416
entomopathogenic nematodes, 416
hazelnut, 416
nematode efficacy, 416
organophosphate, 482, 823, 869, 1400
bioassay, 482
Chrsoperla carnea, 823
citrus, 1400
Cydia pomonella, 482
detoxifying enzymes, 482
dose–mortality analysis, 869
fitness benefit, 823
insecticide resistance, 1400
neonicotinoids, 869
pyrethroids, 823, 869
redbanded stink bug, 869
resistance, 823
resistance detection, 482
Ostrinia nubilalis, 249, 331, 2080
dispersal, 331
ecotype, 249
European corn borer, 2080
field corn, 2080
infestation technique, 2080
parasitoid, 249
plant abandonment, 331
silking behavior, 331
synchronization, 249
transgenic corn, 331
oviposition, 1180, 1744
developmental period, 1744
feeding, 1744
Frankliniella occidentalis, 1744
host plant selection, 1180
mechanism, 1180
preference, 1180
resistance, 1180
survival rate, 1744
oviposition deterrent,
antifeedant, 1196
plant–insect–chemistry triad, 1196
sublethal effects, 1196

P

P450 resistance,
fungicide, 1810
Listronotus maculicollis, 1810
plant growth regulators, 1810
synergist, 1810
palm,
Inesfly IGR FITO, 402
Phoenix conariensis, 402
Rhynchophorus ferrugineus, 402
Panonychus citri,
genetic differentiation, 2204
mitochondrial cytochrome oxidase subunit I, 2204
nucleotide diversity, 2204
population structure, 2204
paraffin wax,
Grapholita molesta, 1299
mating disruption, 1299
pheromone dispenser, 1299
point source, 1299
parasitic mite,
honey bee, 1039
pest management, 1039
parasitoid, 249, 583, 1720, 1770, 1848
Acrobasis vacinii, 1720
Aprostocetus hagenowii, 1770
Bemisia tabaci, 1848
biocontrol, 583
ecotype, 249
freeze, 1770
heat, 1770
infection frequency, 1848
life-stage activity, 1720
Ostrinia nubilalis, 249
Periplaneta americana, 1770
predator, 1848
predatory flies, 583
reduced-risk insecticide, 1720
soybean aphid, 583
synchronization, 249
Wolbachia supergroups, 1848
particle film technology,
apple maggot, 394
behavior, 394
kaolin, 394
Surround WP, 394
path,
dispersal, 63
movement, 63
tracking, 63
pathogens,
Apis mellifera, 1517
classification and regression tree analysis, 1517
colony collapse disorder, 1517
epidemiology, 1517
pathway,
exotic, 1682
firewood, 1682
fuelwood, 1682
invasive, 1682
PBO,
acaricide resistance, 1360
bioassay, 1360
synergist, 1360
verbutin, 1360
pear,
ITS, 25
mealybug, 25
molecular identification, 25
species-specific primer, 25
pear psylla,
abamectin susceptibility, 816
bioassays, 816
insecticide resistance, 816
pest control, 816
Pectinophora gossypiella,
ecological negative cross-resistance, 1821
resistance management, 1821
simulation modeling, 1821
Steinernema riobrave, 1821
Periplaneta americana,
Aprostocetus hagenowii, 1770
freeze, 1770
heat, 1770
parasitoid, 1770
peritrophic membrane,
baiting, 423
chitin synthesis inhibitor, 423
Coptotermes, 423
nest temperature, 423
Persea americana,
Apis mellifera, 228
perseitol, 228
pollination, 228
perseitol,
Apis mellifera, 228
Persea americana, 228
pollination, 228
pest control, 166, 497, 816
abamectin susceptibility, 816
avidin, 497
bioassays, 816
biotin, 497
insecticide resistance, 816
Loxosceles, 166
loxoscelism, 166
pear psylla, 816
pyrethroids, 166
Sicariidae, 166
streptavidin, 497
transgenic plant, 497
pest management, 898, 1039, 1790
Coccidae, 1790
cytochrome oxidase subunit II, 898
honey bee, 1039
Linepithema humile, 1790
Mesolecanium nigrofasciatum, 1790
molecular diagnostic marker, 898
parasitic mite, 1039
red maple, 1790
weevil, 898
pest management tools,
intraguild predation, 1086
pollen, 1086
predators, 1086
western flower thrips, 1086
pest quarantines,
emerald ash borer, 597
firewood, 597
phytosanitation, 597
regulatory compliance, 597
pesticide,
complementary log-log modeling, 1665
rice insect pest, 1665
virulence, 1665
pesticide residues,
Apis mellifera, 1964
colony losses, 1964
stored pollen, 1964
Phaseolus vulgaris,
Aphis glycines, 1670
nonpersistent virus, 1670
seasonal phenology, 1670
phenology,
bark beetles, 1187
exotic species, 1187
Ohio, 1187
semiochemicals, 1187

pheromone, 284, 641
Cerambycidae, 641
Fluon, 641
Pholetesor, 284
population density, 284
trap, 641
trapping, 284
Vaccinium corymbosum, 284
wood-boring insect, 641
pheromone dispenser,
Grapholita molesta, 1299
mating disruption, 1299
paraffin wax., 1299
point source, 1299
pheromone monitoring,
canola, 654
IPM, 654
Lepidoptera, 654
Plutella xylostella, 654
pheromone trapping,
flour mill, 1002
fumigation, 1002
population dynamics, 1002
Tribolium castaneum, 1002
Phoenix conariensis,
Inesfly IGR FITO, 402
palm, 402
Rhynchophorus ferrugineus, 402
Phoenix dactylifera,
binomial sampling plan, 1474
Ectomyelois ceratoniae, 1474
resampling validation, 1474
seasonal population dynamics, 1474
Pholetesor,
pheromone, 284
population density, 284
trapping, 284
Vaccinium corymbosum, 284
phoresy,
Abgrallaspis aguacatae, 1172
Aspidiotus nerii, 1172
Diaspidiotus perniciosus, 1172
Hemiberlesia lanatiae, 1172
phosphine,
Carposina niponensis, 1988
fumigant, 1988
toxicity, 1988
phosphorus deficiency,
biotic, 516
herbivory, 516
stem mining, 516
water stress, 516
Phthorimaea operculella,
desiccation, 1306
insecticide application timing, 1306
integrated pest management, 1306
irrigation, 1306
Phyllocnistis citrella,
control level, 1711
Diaphorina citri, 1711
imidacloprid, 1711
systemic uptake, 1711
Phyllophaga anxia,
entomopathogenic nematodes, 1979
Fraser fir, 1979
soil-applied insecticides, 1979
physiological model,
Acrobasis nuxvorella, 735
biofix, 735
Carya, 735
heat unit, 735
phytohormones,
Mayetiola destructor, 178
resistance, 178
wheat, 178
phytosanitary treatment,
avocado, 1112
systems approach, 1112
Thaumatotibia leucotreta, 1112
phytosanitation,
emerald ash borer, 597
firewood, 597
pest quarantines, 597
regulatory compliance, 597
Phytoseiidae,
biological control, 2061
Calepitrimerus vitis, 2061
Colomerus vitis, 2061
integrated pest management in viticulture, 2061
Pierce's disease,
alternate host plants, 1051
citrus, 1051
grape, 1051
Pierce's disease, 1051
vector control, 1051
Pinus ponderosa,
Dendroctonus, 1693
Ips pini, 1693
microclimate, 1693
thinning, 1693
Pinus yunnanensis,
aggregation, 95
flight pattern, 95
Tomicus yunnanensis, 95
Pisum sativum,
population dynamics, 691
reduced tillage, 691
Sitona lineatus, 691
Planococcus citri,
citrus mealybug, 619
grapevine, 619
Grapevine leafroll-associated virus 3, 619
insect, 619
Planococcus ficus,
controlled atmosphere, 272
grape benchgraft, 272
mealybug, 272
ultralow oxygen, 272
plant abandonment,
dispersal, 331
Ostrinia nubilalis, 331
silking behavior, 331
transgenic corn, 331
plant defense,
artificial infestation, 1866
Brassica, 1866
glucosinolates, 1866
Mamestra brassicae, 1866
plant disease vector,
dispersal, 1524
psyllid yellows, 1524
zebra chip, 1524
plant growth regulators,
fungicide, 1810
Listronotus maculicollis, 1810
P450 resistance, 1810
synergist, 1810
plant pathogen,
carrot, 1060
liberibacter, 1060
psyllid, 1060
Trioza apicalis, 1060
plant resistance, 1412, 2229
boring behavior, 1412
Chilo suppressalis, 1412
Hessian fly genotypes, 2229
Mayetiola destructor, 2229
rice, 1412
silicon, 1412
wheat breeding, 2229
plant–insect–chemistry triad,
antifeedant, 1196
oviposition deterrent, 1196
plant–insect–chemistry triad, 1196
sublethal effects, 1196
Platynota idaeusalis,
apple, 1657
conservation biological control, 1657
extrafloral nectar, 1657
Goniozus floridanus, 1657
Plodia interpunctella,
simulation model, 1503
stored-product insect, 1503
plum curculio,
benzaldehyde, 744
Circle trap, 744
grandisoic acid, 744
pyramid trap, 744
Plutella xylostella, 654, 843, 1347, 2019, 2028
baseline susceptibility, 843
canola, 654
canola cultivars, 2019
chlorantraniliprole, 843
discriminating dose, 843
emamectin benzoate, 1347
emergence period, 2028
emergence rate, 2028
epigenetic transmission, 1347
humidity, 2028
IPM, 654
Lepidoptera, 654
life-table analysis, 2019
model based survival, 2019
pheromone monitoring, 654
resistance, 1347
resistance monitoring, 843
temperature, 2028
tolerance, 1347
point source,
Grapholita molesta, 1299
mating disruption, 1299
paraffin wax., 1299
pheromone dispenser, 1299
pollen,
intraguild predation, 1086
pest management tools, 1086
predators, 1086
western flower thrips, 1086

pollination, 228, 1286
Amegilla, 1286
Apis mellifera, 228
flavor, 1286
horticulture, 1286
Persea americana, 228
perseitol, 228
seeds, 1286
population, 127, 2126
citrus greening, 127
feeding stimulant, 2126
foraging, 2126
Huanglongbing, 127
king mandarin, 127
Reticulitermes flavipes, 2126
sugar, 2126
systemic insecticide, 127
population density,
pheromone, 284
Pholetesor, 284
trapping, 284
Vaccinium corymbosum, 284
population dynamics, 691, 1002, 1729, 2242
Bactericera cockerelli, 1729
flour mill, 1002
fumigation, 1002
global sampling, 2242
life table, 1729
mortality, 1729
mites, 2242
pheromone trapping, 1002
Pisum sativum, 691
powdery mildew, 2242
reduced tillage, 691
Sitona lineatus, 691
thrips, 2242
Tribolium castaneum, 1002
Zebra Chip, 1729
population estimation,
sequential sampling, 525
Taylor's power law, 525
treatment threshold, 525
population growth,
developmental rates, 1920
stored grain, 1920
stored products, 1920
population growth parameter,
demography, 966
potato cultivar, 966
potato tuberworm, 966
Solanum tuberosum, 966
population structure,
genetic differentiation, 2204
mitochondrial cytochrome oxidase subunit I, 2204
nucleotide diversity, 2204
Panonychus citri, 2204
Populus, 718, 2223
biomass production, 718
Chaitophorus leucomelas, 2223
life table, 2223
Lymantria dispar, 718
microbial pesticides, 718
short rotation trees, 718
susceptibility, 2223
postemergence longevity,
emergence, 234
low-temperature tolerance, 234
stress, 234
postharvest,
CATTS, 265
controlled atmosphere, 265
false codling moth, 265
heat, 265
postpharyngeal gland,
ant bait, 676
Malpighian tubules, 676
microscopy, 676
midgut, 676
potato cultivar,
demography, 966
population growth parameter, 966
potato tuberworm, 966
Solanum tuberosum, 966
potato tuberworm,
demography, 966
population growth parameter, 966
potato cultivar, 966
Solanum tuberosum, 966
powdery mildew,
global sampling, 2242
mites, 2242
population dynamics, 2242
thrips, 2242
predation,
Aphis glycines, 1483
field experiments, 1483
host plant resistance, 1483
sampling, 1483
predator, 1086, 1149, 1848
Bemisia tabaci, 1848
biocontrol, 1149
Bradysia, 1149
Coenosia attenuata, 1149
development, 1149
infection frequency, 1848
intraguild predation, 1086
parasitoid, 1848
pest management tools, 1086
pollen, 1086
western flower thrips, 1086
Wolbachia supergroups, 1848
predatory flies,
biocontrol, 583
parasitoids, 583
soybean aphid, 583
predictive model,
Cylindrocopturus adspersus, 302
degree-day model, 302
sunflower stem weevil, 302
preference,
host plant selection, 1180
mechanism, 1180
oviposition, 1180
resistance, 1180
prey-mediated effects,
Laricobius nigrinus, 1222
nontarget, 1222
Sasajiscymnus tsugae, 1222
Tsuga canadensis, 1222
prism trap,
emerald ash borer, 1235
trap height, 1235
visual cues, 1235
product moisture,
Indianmeal moth, 612
relative humidity, 612
vacuum treatment, 612
prophylactic treatment,
bamboo, 101
cost/benefit, 101
Dinoderus minutus, 101
wood preservatives, 101
psyllid,
carrot, 1060
liberibacter, 1060
plant pathogen, 1060
Trioza apicalis, 1060
psyllid yellows,
dispersal, 1524
plant disease vector, 1524
zebra chip, 1524
Pyralidae,
Amyelois transitella, 314
Pyralis farinalis, 314
sex pheromone, 314
Pyralis farinalis,
Amyelois transitella, 314
Pyralidae, 314
sex pheromone, 314
pyramid trap,
benzaldehyde, 744
Circle trap, 744
grandisoic acid, 744
plum curculio, 744
pyrethroid, 166, 823, 869
Chrsoperla carnea, 823
dose–mortality analysis, 869
fitness benefit, 823
Loxosceles, 166
loxoscelism, 166
neonicotinoids, 869
organophosphates, 823, 869
pest control, 166
redbanded stink bug, 869
resistance, 823
Sicariidae, 166
pyrethroid cross-resistance,
Alphitobius diaperinus, 2155
Australia, 2155
darkling beetle, 2155
topical application, 2155
pyriproxyfen,
growth and development, 340
insect growth regulator, 340
lethal and sublethal effects, 340
obliquebanded leafroller, 340

Q

Q biotype,
Bemisia tabaci, 2174
insect growth regulators, 2174
insecticide resistance, 2174
neonicotinoids, 2174
quarantine, 34, 603, 1129, 1950
Agrilus, 603
Anastrepha ludens, 34, 1129
commodity treatment, 34, 1129, 1950
controlled atmosphere, 34
disinfestation, 34, 1129, 1950
Fraxinus, 603
fumigation, 603
irradiation, 1950
radiation, 1129
sulfuryl fluoride, 603

quarantine threat,
Curculio caryae, 1312
new host association, 1312
weevil DNA, 1312

R

rabbiteye,
blueberries, 1258
Dasineura oxycoccana, 1258
distribution, 1258
emergence trap, 1258
radiation,
Anastrepha ludens, 1129
commodity treatment, 1129
disinfestation, 1129
quarantine, 1129
random amplified polymorphic DNA polymerase chain reaction,
abamectin, 2197
cyromazine, 2197
Liriomyza trifolii, 2197
resistance, 2197
rearing methods,
Blissus insularis, 726
chinch bug, 726
insect diet, 726
red flour beetle,
bed bug, 1636
cockroach, 1636
drugstore beetle, 1636
rice weevil, 1636
red maple,
Coccidae, 1790
Linepithema humile, 1790
Mesolecanium nigrofasciatum, 1790
pest management, 1790
redbanded stink bug,
dose–mortality analysis, 869
neonicotinoids, 869
organophosphates, 869
pyrethroid, 869
reduced tillage,
Pisum sativum, 691
population dynamics, 691
Sitona lineatus, 691
reduced-risk insecticide,
Acrobasis vaciniì, 1720
life-stage activity, 1720
parasitoid, 1720
refuge strategy,
resistance management, 849
toxin pyramid, 849
western corn rootworm, 849
refuge-in-a-bag,
Bt corn, 147
resistance management, 147
seed mix refuge, 147
western corn rootworm, 147
regulatory compliance,
emerald ash borer, 597
firewood, 597
pest quarantines, 597
phytosanitation, 597
relative humidity, 612, 1613
Indianmeal moth, 612
landscape, 1613
Leptinotarsa decemlineata, 1613
product moisture, 612
solar radiation, 1613
temperature, 1613
vacuum treatment, 612
repellency, 2118, 2132
Blattella germanica, 2118
Coptotermes formosanus, 2132
food consumption, 2132
formulation, 2118
microencapsulation, 2118
naphthalene derivatives, 2132
toxicity, 2118
reproduction,
Coffea arabica, 1159
Hypothenemus hampei, 1159
infestation, 1159
temperature, 1159
reproductive biology,
insecticide resistance management, 861
sperm precedence, 861
tobacco budworm, 861
YHD2, 861
resampling validation,
binomial sampling plan, 1474
Ectomyelois ceratoniae, 1474
Phoenix dactylifera, 1474
seasonal population dynamics, 1474
residues,
chlorantraniliprole, 1277
Choristoneura rosaceana, 1277
exposure period, 1277
spinetoram, 1277
resistance, 178, 823, 938, 1166, 1180, 1347, 1378, 2052, 2147, 2197
abamectin, 2197
Bacillus thuringiensis, 2147
barley, 938
biotypes, 938
chlorantraniliprole, 1378
Chrsoperla carnea, 823
Cnaphalocrocis medinalis, 1166
Cry1Ac, 2147
Cry2Ab, 2147
cyromazine, 2197
Diabrotica virgifera virgifera, 2052
Diuraphis noxia, 938
emamectin benzoate, 1347
epigenetic transmission, 1347
fitness benefit, 823
Helicoverpa punctigera, 2147
heritability, 1378
host plant selection, 1180
insect behavior, 2052
larvae, 2052
Liriomyza trifolii, 2197
Mayetiola destructor, 178
mechanism, 1180
obliquebanded leafroller, 1378
organophosphate, 823
oviposition, 1180
phytohormones, 178
Plutella xylostella, 1347
preference, 1180
pyrethroids, 823
random amplified polymorphic DNA polymerase chain reaction, 2197
rice variety, 1166
RWA, 938
spinetoram, 1378
tolerance, 1347
wheat, 178
Zea mays, 2052
resistance and insect resistance management,
Cry1Ab, 1386
fitness costs, 1386
high-dose refuge, 1386
resistance detection,
bioassay, 482
Cydia pomonella, 482
detoxifying enzymes, 482
organophosphates, 482
resistance management, 147, 849, 1803, 1821, 2187
Bacillus thuringiensis, 2187
Bt, 2187
Bt corn, 147
Bt maize, 1803
Diabrotica virgifera virgifera, 2187
ecological negative cross-resistance, 1821
European corn borer, 1803
HDR strategy, 1803
mCry3A, 2187
Mon810, 1803
Pectinophora gossypiella, 1821
refuge strategy, 849
refuge-in-a-bag, 147
seed mix refuge, 147
simulation modeling, 1821
Steinernema riobrave, 1821
toxin pyramid, 849
western corn rootworm, 147, 849
resistance mechanism,
Blattella germanica, 460
insecticide resistance, 460
Singapore, 460
synergism, 460
resistance monitoring,
baseline susceptibility, 843
chlorantraniliprole, 843
discriminating dose, 843
Plutella xylostella, 843
resistance to pyriproxyfen,
B biotype, 158
Bemisia tabaci, 158
cytochrome P450, 158
glutathione S-transferase, 158
monooxygenase, 158
responsiveness,
codlemone, 1704
Cydia pomonella, 1704
methoxyfenozide, 1704
sublethal effects, 1704
Reticulitermes flavipes, 762, 791, 2126
colony elimination, 791
colony suppression, 791
feeding stimulant, 2126
foraging, 2126
indoxacarb, 762
microsatellites, 791
population, 2126
Reticulitermes virginicus, 791
sublethal behaviors, 762
sugar, 2126
Reticulitermes virginicus,
colony elimination, 791
colony suppression, 791
microsatellites, 791
Reticulitermes flavipes, 791

Rhagoletis indifferens,
acetamiprid, 379
egg laying, 379
spinosad bait, 379
thiamethoxam, 379
Rhipicephalus microplus,
acaricide resistance, 1355
dieldrin, 1355
GABA receptor, 1355
Rhipicephalus,
Boophilus annulatus, 211
Boophilus microplus, 211
cattle fever tick, 211
white-tailed deer, 211
Rhynchophorus ferrugineus,
Inesfly IGR FITO, 402
palm, 402
Phoenix conariensis, 402
Rhyzopertha dominica, 197, 977, 1508
efficacy assessment, 1508
Heterorhabditis bacteriophora, 977
infrared radiation, 1508
Lepinotus reticulatus, 197
nonchemical method, 1508
Sitophilus oryzae, 197
Steinernema carpocapsae, 977
Steinernema feltiae, 977
stored product pests *kuehniella*, 977
stored products, 197
Tribolium castaneum, 197
rice,
boring behavior, 1412
Chilo suppressalis, 1412
plant resistance, 1412
silicon, 1412
rice insect pest,
complementary log-log modeling, 1665
pesticide, 1665
virulence, 1665
rice variety,
Cnaphalocrocis medinalis, 1166
resistance, 1166
rice weevil,
bed bug, 1636
cockroach, 1636
drugstore beetle, 1636
red flour beetle, 1636
riparian area,
Aphthona, 242
biological control, 242
inundative release, 242
leafy spurge, 242
risk analysis, 1100, 2000
Anastrepha ludens, 2000
Ceratitis capitata, 1100
Ceratitis cosyra, 1100
Ceratitis rosa, 1100
China, 1100
cold treatment, 2000
Hass avocado, 2000
Tephritidae, 2000
RWA,
barley, 938
biotypes, 938
Diuraphis noxia, 938
resistance, 938

S

Saccharum resistance,
intrinsic rate of increase, 1431
Melanaphis sacchari, 1431
Sipha flava, 1431
sampling, 1483, 2236
Aphis glycines, 1483
boll damage, 2236
cotton, 2236
external lesions, 2236
field experiments, 1483
host plant resistance, 1483
predation, 1483
stink bug, 2236
sampling procedure,
Aphis glycines, 186
migration, 186
soybean aphid, 186
suction trap, 186
sanitation,
bagging, 1577
chemical control, 1577
systems approach, 1577
Sasajiscymnus tsugae,
Laricobius nigrinus, 1222
nontarget, 1222
prey-mediated effects, 1222
Tsuga canadensis, 1222
Scarabaeidae,
Anomala orientalis, 1547
Bacillaceae, 1547
Bacillus thuringiensis japonensis, 1547
Cyclocephala borealis, 1547
Schizaphis graminum,
biotype, 1454
genetic diversity, 1454
microsatellite, 1454
simple sequence repeat, 1454
28SD2,
barcoding, 877
cytochrome c oxidase I gene, 877
molecular identification, 877
tospovirus vector, 877
seaborne,
Coptotermes formosanus, 1338
Coptotermes gestroi, 1338
spatial analysis, 1338
termite infestation, 1338
seasonal phenology,
Aphis glycines, 1670
nonpersistent virus, 1670
Phaseolus vulgaris, 1670
seasonal population dynamics,
binomial sampling plan, 1474
Ectomyelois ceratoniae, 1474
Phoenix dactylifera, 1474
resampling validation, 1474
seasonal production,
Bollgard II, 1206
chilling day-degrees, 1206
critical period, 1206
temperature threshold, 1206
seed mix refuge,
Bt corn, 147
refuge-in-a-bag, 147
resistance management, 147
western corn rootworm, 147
seeds,
Amegilla, 1286
flavor, 1286
horticulture, 1286
pollination, 1286
semiochemicals,
bark beetles, 1187
exotic species, 1187
Ohio, 1187
phenology, 1187
SeMNPV,
artificial diet, 17
growth and reproduction, 17
OB yield, 17
Spodoptera exigua, 17
sequential sampling,
population estimation, 525
Taylor's power law, 525
treatment threshold, 525
sericophily,
Arachnida, 808
brown recluse spider, 808
Loxosceles, 808
urban entomology, 808
Sesamia inferens,
Bacillus thuringiensis, 1444
Chilo suppressalis, 1444
Cry1Ab/Vip3H protein, 1444
transgenic rice, 1444
sex pheromone,
Amyelois transitella, 314
Pyralidae, 314
Pyralis farinalis, 314
sexual performance,
Caribbean fruit fly, 40
hydrolyzed yeast, 40
male competitiveness, 40
methoprene, 40
short rotation trees,
biomass production, 718
Lymantria dispar, 718
microbial pesticides, 718
Populus, 718
Sicariidae,
Loxosceles, 166
loxoscelism, 166
pest control, 166
pyrethroids, 166
silicon,
boring behavior, 1412
Chilo suppressalis, 1412
plant resistance, 1412
rice, 1412
silking behavior,
dispersal, 331
Ostrinia nubilalis, 331
plant abandonment, 331
transgenic corn, 331
simple sequence repeat,
biotype, 1454
genetic diversity, 1454
microsatellite, 1454
Schizaphis graminum, 1454
simulation model,
Plodia interpunctella, 1503
stored-product insect, 1503
simulation modeling,
ecological negative cross-resistance, 1821
Pectinophora gossypiella, 1821

resistance management, 1821
Steinernema riobrave, 1821
Singapore,
Blattella germanica, 460
insecticide resistance, 460
resistance mechanism, 460
synergism, 460
single-nucleotide polymorphism,
genetic stock identification, 917
honey bee, 917
microsatellite, 917
single-nucleotide polymorphism-polymerase chain reaction,
B. carambolae, 1994
Bactrocera payayae, 1994
cytochrome oxidase I, 1994
species differentiation, 1994
Sipha flava,
intrinsic rate of increase, 1431
Melanaphis sacchari, 1431
Saccharum resistance, 1431
Sitona lineatus,
Pisum sativum, 691
population dynamics, 691
reduced tillage, 691
Sitophilus oryzae,
Lepinotus reticulatus, 197
Rhyzopertha dominica, 197
stored products, 197
Tribolium castaneum, 197
soil barrier,
dust, 2140
economic cost, 2140
fumigant, 2140
termiticide, 2140
soil penetration,
chemical barrier, 433
fipronil, 433
imidacloprid, 433
termites, 433
soil-applied insecticides,
entomopathogenic nematodes, 1979
Fraser fir, 1979
Phyllophaga anxia, 1979
Solanaceae,
Africa, 70
Bactrocera latifrons, 70
host preference, 70
host range, 70
Solanum tuberosum,
demography, 966
population growth parameter, 966
potato cultivar, 966
potato tuberworm, 966
solar radiation,
landscape, 1613
Leptinotarsa decemlineata, 1613
relative humidity, 1613
temperature, 1613
Solenopsis invicta,
barrier, 754
horizontal toxicity, 754
insecticide, 754
solid phase microextraction,
biological control, 1094
defensive pheromone, 1094
entomopathogenic fungi, 1094
stored products pests, 1094
soybean,
aphid, 509
aphid resistance, 509
biotype, 509
soybean aphid, 509
soybean aphid, 186, 509, 583, 648, 949
aphid, 509
aphid resistance, 509
Aphis glycines, 186, 648
biocontrol, 583
biotype, 509
biotypes, 949
detached leaves, 949
Dowling, 648
electrical penetration graph, 648
host plant resistance, 648
migration, 186
parasitoids, 583
predatory flies, 583
sampling procedure, 186
soybean, 509
soybean host plant resistance, 949
suction trap, 186
soybean cultivars,
demographic parameters, 1420
Helicoverpa armigera, 1420
nutritional indices, 1420
soybean herbivory,
crop rotation, 1464
image analysis, 1464
western corn rootworm, 1464
soybean host plant resistance,
biotypes, 949
detached leaves, 949
soybean aphid, 949
spatial analysis, 533, 1338
Caribbean fruit fly, 533
Coptotermes formosanus, 1338
Coptotermes gestroi, 1338
seaborne, 1338
synthetic lures, 533
termite infestation, 1338
torula yeast, 533
trapping, 533
spatial scale,
attractive range, 1242
codling moth, 1242
mark–release–recapture, 1242
mating disruption, 1242
spatial statistics,
contour analysis, 1886
geostatistics, 1886
Mediterranean fruit fly, 1886
variogram, 1886
species differentiation,
B. carambolae, 1994
Bactrocera payayae, 1994
cytochrome oxidase I, 1994
single-nucleotide polymorphism-polymerase chain reaction, 1994
species richness,
canopy malaise, 698
exotic species survey, 698
trapping, 698
species-specific primer,
ITS, 25
mealybug, 25
molecular identification, 25
pear, 25
sperm precedence,
insecticide resistance management, 861
reproductive biology, 861
tobacco budworm, 861
YHD2, 861
spider mites,
BROTE, 348
downy brome, 348
Neoseiulus, 348
western corn rootworm, 348
spinetoram, 140, 1277, 1378, 1815
acetamiprid, 1815
chlorantraniliprole, 140, 1277, 1378, 1815
Choristoneura rosaceana, 1277
exposure period, 1277
field-collected populations, 140
Grapholita molesta, 1815
heritability, 1378
insecticide resistance, 140
obliquebanded leafroller, 140, 1378
residues, 1277
resistance, 1378
spinosad, 1815
spinosad, 203, 409, 550, 1815
acetamiprid, 1815
Bactrocera cucurbitae, 409
Bactrocera dorsalis, 409
biopesticide, 203
Callosobruchus maculatus, 203
chlorantraniliprole, 1815
cowpea storage, 203
Grapholita molesta, 1815
house fly, 550
imidacloprid, 550
metaflumizone, 550
Musca domestica, 550
naled, 409
spinetoram, 1815
tephritid flies, 409
spinosad bait,
acetamiprid, 379
egg laying, 379
Rhagoletis indifferens, 379
thiamethoxam, 379
spittlebugs,
adult damage, 1860
Brachiaria, 1860
host plant resistance, 1860
life stages, 1860
SPME,
4-8-dimethyldecanal, 1915
aggregation pheromone, 1915
stored products, 1915
Spodoptera exigua, 17, 577, 662
artificial diet, 17
cannibalism, 577
development, 662
fecundity, 662
fertility, 662
growth and reproduction, 17
juvenile hormone, 577
methoxyfenozide, 662
nucleopolyhedrovirus production, 577
OB yield, 17
SeMNPV, 17

Spodoptera frugiperda,
haplotypes, 360
migration, 360
spray coverage, 219, 2107
acetamiprid, 2107
Contarinia nasturtii, 2107
Ethovision, 219
image analysis, 219
insecticide residue, 2107
invasive species, 2107
IPM, 219
water-sensitive spray cards, 219
16S rDNA,
bacterial diversity, 1832
colonization, 1832
ethidium monoazide, 1832
Musca domestica, 1832
stable fly,
Bacillus thuringiensis, 1019
horn fly, 1019
microbial control, 1019
starvation,
Beauveria bassiana, 1542
diet, 1542
stress, 1542
Tribolium castaneum, 1542
Steinernema carpocapsae,
Heterorhabditis bacteriophora, 977
Rhyzopertha dominica, 977
Steinernema feltiae, 977
stored product pests *kuehniella*, 977
Steinernema feltiae,
Heterorhabditis bacteriophora, 977
Rhyzopertha dominica, 977
Steinernema carpocapsae, 977
stored product pests *kuehniella*, 977
Steinernema riobrave,
ecological negative cross-resistance, 1821
Pectinophora gossypiella, 1821
resistance management, 1821
simulation modeling, 1821
Steinernema spp.,
Aethina tumida, 1
Apis mellifera, 1
entomopathogenic nematodes, 1
Heterorhabditis spp., 1
stem mining,
biotic, 516
herbivory, 516
phosphorus deficiency, 516
water stress, 516
sterile insect technique (SIT), 1144, 2034
Coleoptera, 1144
Cylas formicarius elegantulus, 1144, 2034
gamma radiation, 2034
laboratory adaptation, 1144
mating behavior, 2034
substerilization, 2034
sticky trap,
inflorescence, 631
Mangifera indica, 631
nondestructive sampling methods, 631
Thripidae, 631
stink bug,
boll damage, 2236
cotton, 2236
external lesions, 2236
sampling, 2236
stochastic dynamic programming,
insect management, 1896
IPM, 1896
stored corn, 1896
Stomoxys calcitrans,
gene flow, 1012
genetics, 1012
isozyme, 1012
Thailand, 1012
storage mite pests,
diatomaceous earth, 190
stored wheat, 190
Tyrophagus putrescentiae, 190
stored corn,
insect management, 1896
IPM, 1896
stochastic dynamic programming, 1896
stored grain,
developmental rates, 1920
population growth, 1920
stored products, 1920
stored pollen,
Apis mellifera, 1964
colony losses, 1964
pesticide residues, 1964
stored product pests *kuehniella*,
Heterorhabditis bacteriophora, 977
Rhyzopertha dominica, 977
Steinernema carpocapsae, 977
Steinernema feltiae, 977
stored products, 197, 1915, 1920
aggregation pheromone, 1915
developmental rates, 1920
4-8-dimethyldecanal, 1915
Lepinotus reticulatus, 197
population growth, 1920
Rhyzopertha dominica, 197
Sitophilus oryzae, 197
SPME, 1915
stored grain, 1920
Tribolium castaneum, 197
stored products pests,
biological control, 1094
defensive pheromone, 1094
entomopathogenic fungi, 1094
solid phase microextraction, 1094
stored wheat,
diatomaceous earth, 190
storage mite pests, 190
Tyrophagus putrescentiae, 190
stored-product insect,
Plodia interpunctella, 1503
simulation model, 1503
streptavidin,
avidin, 497
biotin, 497
pest control, 497
transgenic plant, 497
stress, 234, 1542
Beauveria bassiana, 1542
diet, 1542
emergence, 234
low-temperature tolerance, 234
postemergence longevity, 234
starvation, 1542
Tribolium castaneum, 1542
Striacosta albicosta,
Cry1Ab, 54
intraguild competition, 54
MON810, 54
YieldGard, 54
sublethal behaviors,
indoxacarb, 762
Reticulitermes flavipes, 762
sublethal effects, 308, 1196, 1704
antifeedant, 1196
biological control, 308
codlemone, 1704
Cydia pomonella, 1704
insecticide, 308
methoxyfenozide, 1704
oviposition deterrent, 1196
plant–insect–chemistry triad, 1196
responsiveness, 1704
Trichogramma evanescens, 308
Trichogramma semblidis, 308
substerilization,
Cylas formicarius elegantulus, 2034
gamma radiation, 2034
mating behavior, 2034
sterile insect technique (SIT), 2034
subterranean termites,
benzoyl urea insect growth regulator, 770
chitin synthesis inhibitor, 770
termite baits, 770
termite colony elimination, 770
suction trap,
Aphis glycines, 186
migration, 186
sampling procedure, 186
soybean aphid, 186
sugar, 1647, 2126
carbohydrate, 1647
Coptotermes formosanus, 1647
2-deoxy-d-galactose, 1647
feeding stimulant, 2126
foraging, 2126
population, 2126
Reticulitermes flavipes, 2126
termite, 1647
sulfuryl fluoride, 277, 603
Agrilus, 603
alternative fumigation, 277
Bambusa, 277
Chlorophorus annularis, 277
Fraxinus, 603
fumigation, 603
quarantine, 603
sunflower stem weevil,
Cylindrocopturus adspersus, 302
degree-day model, 302
predictive model, 302
supercooling point,
Anobiidae, 1909
cold, 1909
heat, 1909
tolerance, 1909
Surround WP,
apple maggot, 394
behavior, 394
kaolin, 394
particle film technology, 394

surveillance,
invasive fruit flies, 2042
male lures, 2042
mark–recapture, 2042
survival, 1250, 1754
agonism, 1754
almonds, 1250
degree-days, 1250
development, 1250
interspecific, 1754
intraspecific, 1754
Microcerotermes crassus, 1754
navel orangeworm, 1250
survival rate,
developmental period, 1744
feeding, 1744
Frankliniella occidentalis, 1744
oviposition, 1744
survivorship,
alfalfa weevil, 292
fecundity, 292
life table entropy, 292
modeling, 292
susceptibility, 374, 2223
aphids, 374
Aphis pomi, 374
Aphis spiraecola, 374
Chaitophorus leucomelas, 2223
insecticide resistance, 374
life table, 2223
Populus, 2223
sweetpotato,
Conoderus vespertinus, 1268
tobacco wireworm, 1268
wireworm, 1268
Symploce pallens,
fecundity, 443
longevity, 443
nymphal development, 443
synchronization,
ecotype, 249
Ostrinia nubilalis, 249
parasitoid, 249
synergism,
Blattella germanica, 460
insecticide resistance, 460
resistance mechanism, 460
Singapore, 460
synergist, 1360, 1365, 1810
acaricide resistance, 1360
bioassay, 1360
fungicide, 1810
insecticide resistance, 1365
LD-P line, 1365
Leptinotarsa decemlineata, 1365
Listronotus maculicollis, 1810
P450 resistance, 1810
PBO, 1360
plant growth regulators, 1810
topical bioassay, 1365
verbutin, 1360
synthetic lures,
Caribbean fruit fly, 533
spatial analysis, 533
torula yeast, 533
trapping, 533
synthetic trail pheromone,
Argentine ant, 1784
control, 1784
Linepithema humile, 1784
(Z)-9-hexadecenal, 1784
systemic injection,
azadirachtin, 708
emerald ash borer, 708
foliar uptake, 708
larval mortality, 708
systemic insecticide, 127, 1735
armored scales, 1735
bark absorption, 1735
Christmas tree, 1735
citrus greening, 127
Huanglongbing, 127
IPM, 1735
king mandarin, 127
population, 127
systemic uptake,
control level, 1711
Diaphorina citri, 1711
imidacloprid, 1711
Phyllocnistis citrella, 1711
systems approach, 1112, 1577
avocado, 1112
bagging, 1577
chemical control, 1577
phytosanitary treatment, 1112
sanitation, 1577
Thaumatotibia leucotreta, 1112

T

Taylor's power law,
population estimation, 525
sequential sampling, 525
Taylor's power law, 525
treatment threshold, 525
temperature, 1159, 1613, 2028
Coffea arabica, 1159
emergence period, 2028
emergence rate, 2028
humidity, 2028
Hypothenemus hampei, 1159
infestation, 1159
landscape, 1613
Leptinotarsa decemlineata, 1613
Plutella xylostella, 2028
relative humidity, 1613
reproduction, 1159
solar radiation, 1613
temperature threshold,
Bollgard II, 1206
chilling day-degrees, 1206
critical period, 1206
seasonal production, 1206
tephritid flies,
Bactrocera cucurbitae, 409
Bactrocera dorsalis, 409
naled, 409
spinosad, 409
Tephritidae,
Anastrepha ludens, 2000
cold treatment, 2000
Hass avocado, 2000
risk analysis, 2000
termite, 433, 1647
carbohydrate, 1647
chemical barrier, 433
Coptotermes formosanus, 1647
2-deoxy-d-galactose, 1647
fipronil, 433
imidacloprid, 433
soil penetration, 433
sugar, 1647
termite baits,
benzoyl urea insect growth regulator, 770
chitin synthesis inhibitor, 770
subterranean termites, 770
termite colony elimination, 770
termite colony elimination,
benzoyl urea insect growth regulator, 770
chitin synthesis inhibitor, 770
subterranean termites, 770
termite baits, 770
termite infestation,
Coptotermes formosanus, 1338
Coptotermes gestroi, 1338
seaborne, 1338
spatial analysis, 1338
termiticide,
dust, 2140
economic cost, 2140
fumigant, 2140
soil barrier, 2140
Tetranychus urticae,
botanical acaricide, 1293
essential oil, 1293
natural fumigant, 1293
Neoseiulus californicus, 1293
Tetranychus viennensis,
endoplasmic reticulum, 1372
methyl palmitate, 1372
mitochondria, 1372
Thailand,
gene flow, 1012
genetics, 1012
isozyme, 1012
Stomoxys calcitrans, 1012
Thaumatotibia leucotreta,
avocado, 1112
phytosanitary treatment, 1112
systems approach, 1112
thiamethoxam,
acetamiprid, 379
egg laying, 379
Rhagoletis indifferens, 379
spinosad bait, 379
thinning,
Dendroctonus, 1693
Ips pini, 1693
microclimate, 1693
Pinus ponderosa, 1693
Thripidae,
inflorescence, 631
Mangifera indica, 631
nondestructive sampling methods, 631
sticky trap, 631
thrips,
global sampling, 2242
mites, 2242
population dynamics, 2242
powdery mildew, 2242
Thrips tabaci, 925, 1315
action threshold, 1315
Allium cepa, 925, 1315
insecticide, 1315
Iris yellow spot virus, 925
onion resistance, 925

Thysanoptera,
adults, 2164
insecticide resistance susceptibility, 2164
larvae, 2164
tobacco budworm,
insecticide resistance management, 861
reproductive biology, 861
sperm precedence, 861
YHD2, 861
tobacco wireworm,
Conoderus vespertinus, 1268
sweetpotato, 1268
wireworm, 1268
tolerance, 1347, 1405, 1909
Anobiidae, 1909
antibiosis, 1405
Aphis glycines, 1405
cold, 1909
emamectin benzoate, 1347
epigenetic transmission, 1347
Glycine max, 1405
heat, 1909
host plant resistance, 1405
Plutella xylostella, 1347
resistance, 1347
supercooling point, 1909
Tomicus yunnanensis,
aggregation, 95
flight pattern, 95
Pinus yunnanensis, 95
topical application,
Alphitobius diaperinus, 2155
Australia, 2155
darkling beetle, 2155
pyrethroid cross-resistance, 2155
topical bioassay,
insecticide resistance, 1365
LD-P line, 1365
Leptinotarsa decemlineata, 1365
synergist, 1365
torula yeast,
Caribbean fruit fly, 533
spatial analysis, 533
synthetic lures, 533
trapping, 533
tospovirus vector,
barcoding, 877
cytochrome c oxidase I gene, 877
molecular identification, 877
28SD2, 877
toxicity, 448, 1988, 2118
Blattella germanica, 448, 2118
Carposina niponensis, 1988
essential oils, 448
formulation, 2118
fumigant, 1988
microencapsulation, 2118
ootheca, 448
phosphine, 1988
repellency, 2118
toxin pyramid,
refuge strategy, 849
resistance management, 849
western corn rootworm, 849
tracking,
dispersal, 63
movement, 63
path, 63
transgenic corn,
dispersal, 331
Ostrinia nubilalis, 331
plant abandonment, 331
silking behavior, 331
transgenic maize, 1031, 1394
Bt maize, 1394
Cry1Ab, 1394
Cry1F, 1031
Cry3Bb, 1394
fall armyworm, 1031
insect resistance evolution, 1394
transgenic plant,
avidin, 497
biotin, 497
pest control, 497
streptavidin, 497
transgenic rice,
Bacillus thuringiensis, 1444
Chilo suppressalis, 1444
Cry1Ab/Vip3H protein, 1444
Sesamia inferens, 1444
trap,
Cerambycidae, 641
Fluon, 641
pheromone, 641
wood-boring insect, 641
trap height,
emerald ash borer, 1235
prism trap, 1235
visual cues, 1235
trapping, 284, 356, 533, 698
Anthonomus grandis grandis, 356
boll weevil, 356
canopy malaise, 698
Caribbean fruit fly, 533
eradication, 356
exotic species survey, 698
grandlure, 356
pheromone, 284
Pholetesor, 284
population density, 284
spatial analysis, 533
species richness, 698
synthetic lures, 533
torula yeast, 533
Vaccinium corymbosum, 284
treatment threshold,
population estimation, 525
sequential sampling, 525
Taylor's power law, 525
Tribolium castaneum, 197, 991, 1002, 1542
Beauveria bassiana, 1542
diet, 1542
flour mill, 991, 1002
fumigation, 991, 1002
Lepinotus reticulatus, 197
methyl bromide, 991
monitoring, 991
pheromone trapping, 1002
population dynamics, 1002
Rhyzopertha dominica, 197
Sitophilus oryzae, 197
starvation, 1542
stored products, 197
stress, 1542
Trichogramma evanescens,
biological control, 308
insecticide, 308
sublethal effects, 308
Trichogramma semblidis, 308
Trichogramma semblidis,
biological control, 308
insecticide, 308
sublethal effects, 308
Trichogramma evanescens, 308
trimedlure,
attractants, 1586
ceralure, 1586
detection, 1586
Mediterranean fruit fly, 1586
Trioza apicalis,
carrot, 1060
liberibacter, 1060
plant pathogen, 1060
psyllid, 1060
Tsuga canadensis,
Laricobius nigrinus, 1222
nontarget, 1222
prey-mediated effects, 1222
Sasajiscymnus tsugae, 1222
tunneling,
Coptotermes formosanus, 799
desiccation, 799
wood consumption, 799
tunneling activity,
Coptotermes gestroi, 437
Microcerotermes crassus, 437
moisture, 437
wood consumption, 437
twospotted spider mite,
coverage, 386
integrated pest management, 386
mineral oil, 386
Tyrophagus putrescentiae,
diatomaceous earth, 190
storage mite pests, 190
stored wheat, 190

U

ultralow oxygen,
controlled atmosphere, 272
grape benchgraft, 272
mealybug, 272
Planococcus ficus, 272
urban entomology,
Arachnida, 808
brown recluse spider, 808
Loxosceles, 808
sericophily, 808
urban forest,
emerald ash borer, 108
invasive species, 108
nonindigenous species, 108

V

Vaccinium corymbosum, 284, 557, 668
alternative pollinator, 557
Apis mellifera, 557
Apoidea, 668
Bombus impatiens, 557
climate, 557
insecticides, 668
nontarget effects, 668
pheromone, 284
Pholetesor, 284
population density, 284
trapping, 284
wild bee conservation, 668

vacuum treatment,
Indianmeal moth, 612
product moisture, 612
relative humidity, 612
varietal resistance,
Liposcelis, 985
wheat classes, 985
variogram,
contour analysis, 1886
geostatistics, 1886
Mediterranean fruit fly, 1886
spatial statistics, 1886
Varroa destructor, 10, 1797
Aethina tumida, 10
Apis mellifera, 10, 1797
honey bee, 10
monitoring insecticide resistance, 1797
Nosema ceranae, 10
vector,
acquisition, 1531
citrus greening disease, 1531
huanglongbing, 1531
inoculation, 1531
vector control,
alternate host plants, 1051
citrus, 1051
grape, 1051
Pierce's disease, 1051
verbutin,
acaricide resistance, 1360
bioassay, 1360
PBO, 1360
synergist, 1360
Vip3Aa,
cross-tolerance, 2169
Cry1Ac, 2169
Helicoverpa armigera, 2169
viral enhancement,
granulovirus proteins, 257
Helicoverpa armigera, 257
Mamestra brassicae, 257, 257
nucleopolyhedrovirus, 257
virulence,
complementary log-log modeling, 1665
pesticide, 1665
rice insect pest, 1665
visual cues,
emerald ash borer, 1235
prism trap, 1235
trap height, 1235

W

water stress,
biotic, 516
herbivory, 516
phosphorus deficiency, 516
stem mining, 516
water-sensitive spray cards,
Ethovision, 219
image analysis, 219
IPM, 219
spray coverage, 219
weevil,
cytochrome oxidase subunit II, 898
molecular diagnostic marker, 898
pest management, 898
weevil DNA,
Curculio caryae, 1312
new host association, 1312
quarantine threat, 1312
weighted mean model,
foraging territory, 2112
Lincoln index, 2112
mark–release–recapture, 2112
Microcerotermes diversus, 2112
western corn rootworm, 147, 348, 849, 1464
BROTE, 348
Bt corn, 147
crop rotation, 1464
downy brome, 348
image analysis, 1464
Neoseiulus, 348
refuge strategy, 849
refuge-in-a-bag, 147
resistance management, 147, 849
seed mix refuge, 147
soybean herbivory, 1464
spider mites, 348
toxin pyramid, 849
western flower thrips,
intraguild predation, 1086
pest management tools, 1086
pollen, 1086
predators, 1086
western tarnished plant bug,
alternate host, 2009
host suitability, 2009
life table, 2009
Lygus hesperus, 2009
wheat,
Mayetiola destructor, 178
phytohormones, 178
resistance, 178
wheat breeding,
Hessian fly genotypes, 2229
Mayetiola destructor, 2229
plant resistance, 2229
wheat classes,
Liposcelis, 985
varietal resistance, 985
white-tailed deer,
Boophilus annulatus, 211
Boophilus microplus, 211
cattle fever tick, 211
Rhipicephalus, 211
wild bee conservation,
Apoidea, 668
insecticides, 668
nontarget effects, 668
Vaccinium corymbosum, 668
wireworm, 1268, 2087
Conoderus vespertinus, 1268
Conoderus spp., 2087
crop rotation, 2087
Melanotus communis, 2087
sweetpotato, 1268
tobacco wireworm, 1268
Wolbachia supergroups,
Bemisia tabaci, 1848
infection frequency, 1848
parasitoid, 1848
predator, 1848
wood consumption, 437, 799
Coptotermes formosanus, 799
Coptotermes gestroi, 437
desiccation, 799
Microcerotermes crassus, 437
moisture, 437
tunneling, 799
tunneling activity, 437
wood preservatives,
bamboo, 101
cost/benefit, 101
Dinoderus minutus, 101
prophylactic treatment, 101
wood-boring insect,
Cerambycidae, 641
Fluon, 641
pheromone, 641
trap, 641

X

X radiation,
Alpha radiation, 85
Anastrepha fraterculus, 85
Ceratitis capitata, 85
irradiation, 85
Xylosandrus germanus,
conophthorin, 2094
Gnathotrichus materiarius, 2094
Ips grandicollis, 2094
Ips pini, 2094

Y

YHD2,
insecticide resistance management, 861
reproductive biology, 861
sperm precedence, 861
tobacco budworm, 861
YieldGard,
Cry1Ab, 54
intraguild competition, 54
MON810, 54
Striacosta albicosta, 54

Z

Zea mays,
Diabrotica virgifera virgifera, 2052
insect behavior, 2052
larvae, 2052
resistance, 2052
zebra chip, 1524, 1729
Bactericera cockerelli, 1729
dispersal, 1524
life table, 1729
mortality, 1729
plant disease vector, 1524
population dynamics, 1729
psyllid yellows, 1524
Zeuzera pyrina,
insect population, 1621
mating disruption, 1621
olive yield, 1621
(Z)-9-hexadecenal,
Argentine ant, 1784
control, 1784
Linepithema humile, 1784
synthetic trail pheromone, 1784
(Z)-9-tricosene,
house fly, 1929
muscalure, 1929
(3Z6Z9Z12Z15Z)-pentacosapentaene,
Dioryctria amatella, 1216
Dioryctria clarioralis, 1216
Dioryctria disclusa, 1216
Dioryctria merkeli, 1216